

Selected Bibliography: Furniture Flame Retardants, Toxicity and Health

Prepared by Green Science Policy Institute

June 2013

1. Abb, M., Stahl, B., & Lorenz, W. (2011). Analysis of brominated flame retardants in house dust. *Chemosphere*, 85(11), 1657–63. doi:10.1016/j.chemosphere.2011.06.022
2. Adamkiewicz, G., Zota, A. R., Fabian, M. P., Chahine, T., Julien, R., Spengler, J. D., & Levy, J. I. (2011). Moving environmental justice indoors: understanding structural influences on residential exposure patterns in low-income communities. *American journal of public health*, 101 Suppl , S238–45. doi:10.2105/AJPH.2011.300119
3. Akutsu, K., Takatori, S., Nozawa, S., Yoshiike, M., Nakazawa, H., Hayakawa, K., Makino, T., et al. (2008). Polybrominated diphenyl ethers in human serum and sperm quality. *Bulletin of environmental contamination and toxicology*, 80(4), 345–50. doi:10.1007/s00128-008-9370-4
4. Al-Mousa, F., & Michelangeli, F. (2012). Some commonly used brominated flame retardants cause Ca²⁺-ATPase inhibition, beta-amyloid peptide release and apoptosis in SH-SY5Y neuronal cells. *PLoS one*, 7(4), e33059. doi:10.1371/journal.pone.0033059
5. Aleksa, K., Carnevale, A., Goodyer, C., & Koren, G. (2012). Detection of polybrominated biphenyl ethers (PBDEs) in pediatric hair as a tool for determining in utero exposure. *Forensic science international*, 218(1-3), 37–43. doi:10.1016/j.forsciint.2011.10.003
6. Ali, N., Van den Eede, N., Dirtu, A. C., Neels, H., & Covaci, A. (2012). Assessment of human exposure to indoor organic contaminants via dust ingestion in Pakistan. *Indoor air*, 22(3), 200–11. doi:10.1111/j.1600-0668.2011.00757.x
7. Ali, Nadeem, Ali, L., Mehdi, T., Dirtu, A. C., Al-Shammari, F., Neels, H., & Covaci, A. (2013). Levels and profiles of organochlorines and flame retardants in car and house dust from Kuwait and Pakistan: implication for human exposure via dust ingestion. *Environment international*, 55, 62–70. doi:10.1016/j.envint.2013.02.001
8. Ali, Nadeem, Dirtu, A. C., Van den Eede, N., Goosey, E., Harrad, S., Neels, H., 't Mannetje, A., et al. (2012). Occurrence of alternative flame retardants in indoor dust from New Zealand: indoor sources and human exposure assessment. *Chemosphere*, 88(11), 1276–82. doi:10.1016/j.chemosphere.2012.03.100
9. Ali, Nadeem, Harrad, S., Goosey, E., Neels, H., & Covaci, A. (2011). “Novel” brominated flame retardants in Belgian and UK indoor dust: implications for human exposure. *Chemosphere*, 83(10), 1360–5. doi:10.1016/j.chemosphere.2011.02.078
10. Ali, Nadeem, Harrad, S., Muenhor, D., Neels, H., & Covaci, A. (2011). Analytical characteristics and determination of major novel brominated flame retardants (NBFRs) in indoor dust. *Analytical and bioanalytical chemistry*, 400(9), 3073–83. doi:10.1007/s00216-011-4966-7
11. Ali, Nadeem, Malik, R. N., Mehdi, T., Eqani, S. A. M. A. S., Javeed, A., Neels, H., & Covaci, A. (2013). Organohalogenated contaminants (OHCs) in the serum and hair of pet cats and dogs: biosentinels of indoor pollution. *The Science of the total environment*, 449, 29–36. doi:10.1016/j.scitotenv.2013.01.037
12. Allen, J. G., McClean, M. D., Stapleton, H. M., & Webster, T. F. (2008). Linking PBDEs in house dust to consumer products using X-ray fluorescence. *Environmental science & technology*, 42(11), 4222–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18589991>
13. Alm, H., Scholz, B., Kultima, K., Nilsson, A., Andrén, P. E., Savitski, M. M., Bergman, A., et al. (2010). In vitro neurotoxicity of PBDE-99: immediate and concentration-dependent effects on protein expression in cerebral cortex cells. *Journal of proteome research*, 9(3), 1226–35. doi:10.1021/pr900723c
14. An, J., Li, S., Zhong, Y., Wang, Y., Zhen, K., Zhang, X., Wang, Y., et al. (2011). The cytotoxic effects of synthetic 6-hydroxylated and 6-methoxylated polybrominated diphenyl ether 47 (BDE47). *Environmental toxicology*, 26(6), 591–9. doi:10.1002/tox.20582

15. Antignac, J.-P., Cariou, R., Maume, D., Marchand, P., Monteau, F., Zalko, D., Berrebi, A., et al. (2008). Exposure assessment of fetus and newborn to brominated flame retardants in France: preliminary data. *Molecular nutrition & food research*, 52(2), 258–65. doi:10.1002/mnfr.200700077
16. Araki, A., Saito, I., Kanazawa, A., Morimoto, K., Nakayama, K., Shibata, E., Tanaka, M., et al. (2013). Phosphorus flame retardants in indoor dust and their relation to asthma and allergies of inhabitants. *Indoor air*. doi:10.1111/ina.12054
17. Arbuckle, T. E., Kubwabo, C., Walker, M., Davis, K., Lalonde, K., Kosarac, I., Wen, S. W., et al. (2013). Umbilical cord blood levels of perfluoroalkyl acids and polybrominated flame retardants. *International journal of hygiene and environmental health*, 216(2), 184–94. doi:10.1016/j.ijheh.2012.03.004
18. Asante, K. A., Adu-Kumi, S., Nakahiro, K., Takahashi, S., Isobe, T., Sudaryanto, A., Devanathan, G., et al. (2011). Human exposure to PCBs, PBDEs and HBCDs in Ghana: Temporal variation, sources of exposure and estimation of daily intakes by infants. *Environment international*, 37(5), 921–8. doi:10.1016/j.envint.2011.03.011
19. Babrauskas, V., Blum, A., R., D., & L., B. (2011). Flame Retardants in Furniture Foam: Benefits and Risks. *Fire Safety Science*, 10, 265–278. Retrieved from <http://www.iafss.org/publications/fss/10/265/>
20. Banasik, M., & Suchecka, D. (2011). Neurotoxicity of PBDEs on the developing nervous system. *Environmental health perspectives*, 119(8), A331; author reply A331–2. doi:10.1289/ehp.1103907
21. Batterman, S., Godwin, C., Chernyak, S., Jia, C., & Charles, S. (2010). Brominated flame retardants in offices in Michigan, USA. *Environment international*, 36(6), 548–56. doi:10.1016/j.envint.2010.04.008
22. Berr, J. S., Stapleton, H. M., & Mitchelmore, C. L. (2010). Accumulation and DNA damage in fathead minnows (*Pimephales promelas*) exposed to 2 brominated flame-retardant mixtures, Firemaster 550 and Firemaster BZ-54. *Environmental toxicology and chemistry / SETAC*, 29(3), 722–9. doi:10.1002/etc.94
23. Behnisch, P. (2003). Brominated dioxin-like compounds: in vitro assessment in comparison to classical dioxin-like compounds and other polyaromatic compounds. *Environment International*, 29(6), 861–877. doi:10.1016/S0160-4120(03)00105-3
24. Bellés, M., Alonso, V., Linares, V., Albina, M. L., Sirvent, J. J., Domingo, J. L., & Sánchez, D. J. (2010). Behavioral effects and oxidative status in brain regions of adult rats exposed to BDE-99. *Toxicology letters*, 194(1-2), 1–7. doi:10.1016/j.toxlet.2010.01.010
25. Bergh, C., Magnus Åberg, K., Svartengren, M., Emenius, G., & Östman, C. (2011). Organophosphate and phthalate esters in indoor air: a comparison between multi-storey buildings with high and low prevalence of sick building symptoms. *Journal of environmental monitoring*: JEM, 13(7), 2001–9. doi:10.1039/c1em10152h
26. Basis, A., & Samara, C. (2012). Polybrominated diphenyl ethers (PBDEs) in the indoor and outdoor environments--a review on occurrence and human exposure. *Environmental pollution (Barking, Essex*: 1987), 169, 217–29. doi:10.1016/j.envpol.2012.04.009
27. Beth-Hubner, M. (1999). Toxicological evaluation and classification of the genotoxic, carcinogenic, reprotoxic and sensitising potential of tris(2-chloroethyl)phosphate. *International Archives of Occupational and Environmental Health*, 72(0), M017–M023–M023. Retrieved from <http://www.springerlink.com/content/u3rvnmjcc02h1v12/>
28. Betts, K. S. (2010). Endocrine damper? Flame retardants linked to male hormone, sperm count changes. *Environmental health perspectives*, 118(3), A 130. doi:10.1289/ehp.118-a130b
29. Blake, C. A., McCoy, G. L., Hui, Y. Y., & LaVoie, H. A. (2011). Perinatal exposure to low-dose DE-71 increases serum thyroid hormones and gonadal osteopontin gene expression. *Experimental biology and medicine (Maywood, N.J.)*, 236(4), 445–55. doi:10.1258/ebm.2010.010334
30. Boas, M., Feldt-Rasmussen, U., & Main, K. M. (2012). Thyroid effects of endocrine disrupting chemicals. *Molecular and cellular endocrinology*, 355(2), 240–8. doi:10.1016/j.mce.2011.09.005

31. Bondy, G. S., Gaertner, D., Cherry, W., MacLellan, E., Coady, L., Arnold, D. L., Doucet, J., et al. (2011). Brominated diphenyl ether (BDE) levels in liver, adipose, and milk from adult and juvenile rats exposed by gavage to the DE-71 technical mixture. *Environmental toxicology*, 26(6), 677–90. doi:10.1002/tox.20603
32. Bradman, A., Castorina, R., Sjödin, A., Fenster, L., Jones, R. S., Harley, K. G., Chevrier, J., et al. (2012). Factors associated with serum polybrominated diphenyl ether (PBDE) levels among school-age children in the CHAMACOS cohort. *Environmental science & technology*, 46(13), 7373–81. doi:10.1021/es3003487
33. Brommer, S., Harrad, S., Van den Eede, N., & Covaci, A. (2012). Concentrations of organophosphate esters and brominated flame retardants in German indoor dust samples. *Journal of environmental monitoring*: JEM, 14(9), 2482–7. doi:10.1039/c2em30303e
34. Bruchajzer, Elżbieta. (2011). Porphyrogenic effect of pentabromodiphenyl ether after repeated administration to rats. *Archives of toxicology*, 85(8), 965–74. doi:10.1007/s00204-010-0621-x
35. Bruchajzer, Elżbieta, Frydrych, B., Sporny, S., & Szymańska, J. A. (2010). Toxicity of penta- and decabromodiphenyl ethers after repeated administration to rats: a comparative study. *Archives of toxicology*, 84(4), 287–99. doi:10.1007/s00204-009-0495-y
36. Bruchajzer, Elżbieta, Frydrych, B., Sporny, S., & Szymańska, J. A. (2011). The effect of short-term intoxication of rats with pentabromodiphenyl ether (in mixture mimic commercial products). *Human & experimental toxicology*, 30(5), 363–78. doi:10.1177/09603271110371261
37. Butt, C. M., Wang, D., & Stapleton, H. M. (2011). Halogenated phenolic contaminants inhibit the in vitro activity of the thyroid-regulating deiodinases in human liver. *Toxicological sciences*: an official journal of the Society of Toxicology, 124(2), 339–47. doi:10.1093/toxsci/kfr117
38. Buttke, D. E., Wolkin, A., Stapleton, H. M., & Miranda, M. L. (2013). Associations between serum levels of polybrominated diphenyl ether (PBDE) flame retardants and environmental and behavioral factors in pregnant women. *Journal of exposure science & environmental epidemiology*, 23(2), 176–82. doi:10.1038/jes.2012.67
39. Cai, Y., Zhang, W., Hu, J., Sheng, G., Chen, D., & Fu, J. (2011). Characterization of maternal transfer of decabromodiphenyl ether (BDE-209) administered to pregnant Sprague-Dawley rats. *Reproductive toxicology (Elmsford, N.Y.)*, 31(1), 106–10. doi:10.1016/j.reprotox.2010.08.005
40. Carmichael, S. L., Herring, A. H., Sjödin, A., Jones, R., Needham, L., Ma, C., Ding, K., et al. (2010). Hypospadias and halogenated organic pollutant levels in maternal mid-pregnancy serum samples. *Chemosphere*, 80(6), 641–6. doi:10.1016/j.chemosphere.2010.04.055
41. Carpenter, D. O. (2011). Health effects of persistent organic pollutants: the challenge for the Pacific Basin and for the world. *Reviews on environmental health*, 26(1), 61–9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21714383>
42. Casals-Casas, C., & Desvergne, B. (2011). Endocrine disruptors: from endocrine to metabolic disruption. *Annual review of physiology*, 73, 135–62. doi:10.1146/annurev-physiol-012110-142200
43. Casas, M., Chevrier, C., Hond, E. Den, Fernandez, M. F., Pierik, F., Philippat, C., Slama, R., et al. (2013). Exposure to brominated flame retardants, perfluorinated compounds, phthalates and phenols in European birth cohorts: ENRIECO evaluation, first human biomonitoring results, and recommendations. *International journal of hygiene and environmental health*, 216(3), 230–42. doi:10.1016/j.ijheh.2012.05.009
44. Caserta, D., Mantovani, A., Marci, R., Fazi, A., Ciardo, F., La Rocca, C., Maranghi, F., et al. (2011). Environment and women's reproductive health. *Human reproduction update*, 17(3), 418–33. doi:10.1093/humupd/dmq061
45. Chao, H.-R., Wang, S.-L., Lee, W.-J., Wang, Y.-F., & Päpke, O. (2007). Levels of polybrominated diphenyl ethers (PBDEs) in breast milk from central Taiwan and their relation to infant birth outcome and maternal menstruation effects. *Environment international*, 33(2), 239–45. doi:10.1016/j.envint.2006.09.013
46. Chen, A., Chung, E., DeFranco, E. A., Pinney, S. M., & Dietrich, K. N. (2011). Serum PBDEs and age at menarche in adolescent girls: analysis of the National Health and Nutrition Examination Survey 2003-2004. *Environmental research*, 111(6), 831–7. doi:10.1016/j.envres.2011.05.016

47. Chen, A., Park, J.-S., Linderholm, L., Rhee, A., Petreas, M., DeFranco, E. A., Dietrich, K. N., et al. (2013). Hydroxylated polybrominated diphenyl ethers in paired maternal and cord sera. *Environmental science & technology*, 47(8), 3902–8. doi:10.1021/es3046839
48. Chen, L., Yu, K., Huang, C., Yu, L., Zhu, B., Lam, P. K. S., Lam, J. C. W., et al. (2012). Prenatal transfer of polybrominated diphenyl ethers (PBDEs) results in developmental neurotoxicity in zebrafish larvae. *Environmental science & technology*, 46(17), 9727–34. doi:10.1021/es302119g
49. Chen, S.-J., Ma, Y.-J., Wang, J., Chen, D., Luo, X.-J., & Mai, B.-X. (2009). Brominated flame retardants in children's toys: concentration, composition, and children's exposure and risk assessment. *Environmental science & technology*, 43(11), 4200–6. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19569352>
50. Chen, S.-J., Ma, Y.-J., Wang, J., Tian, M., Luo, X.-J., Chen, D., & Mai, B.-X. (2010). Measurement and human exposure assessment of brominated flame retardants in household products from South China. *Journal of hazardous materials*, 176(1-3), 979–84. doi:10.1016/j.jhazmat.2009.11.138
51. Chevrier, J., Harley, K. G., Bradman, A., Gharbi, M., Sjödin, A., & Eskenazi, B. (2010). Polybrominated diphenyl ether (PBDE) flame retardants and thyroid hormone during pregnancy. *Environmental health perspectives*, 118(10), 1444–9. doi:10.1289/ehp.1001905
52. Chevrier, J., Harley, K. G., Bradman, A., Sjödin, A., & Eskenazi, B. (2011). Prenatal exposure to polybrominated diphenyl ether flame retardants and neonatal thyroid-stimulating hormone levels in the CHAMACOS study. *American journal of epidemiology*, 174(10), 1166–74. doi:10.1093/aje/kwr223
53. Choi, J.-W., Fujimaki, T. S., Kitamura, K., Hashimoto, S., Ito, H., Suzuki, N., Sakai, S., et al. (2003). Polybrominated dibenzo-p-dioxins, dibenzofurans, and diphenyl ethers in Japanese human adipose tissue. *Environmental science & technology*, 37(5), 817–21. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12666907>
54. Christen, V., Crettaz, P., Oberli-Schrämli, A., & Fent, K. (2010). Some flame retardants and the antimicrobials triclosan and triclocarban enhance the androgenic activity in vitro. *Chemosphere*, 81(10), 1245–52. doi:10.1016/j.chemosphere.2010.09.031
55. Costa, L. G., & Giordano, G. (2007). Developmental neurotoxicity of polybrominated diphenyl ether (PBDE) flame retardants. *Neurotoxicology*, 28(6), 1047–67. doi:10.1016/j.neuro.2007.08.007
56. Costa, L. G., & Giordano, G. (2011). Is decabromodiphenyl ether (BDE-209) a developmental neurotoxicant? *Neurotoxicology*, 32(1), 9–24. doi:10.1016/j.neuro.2010.12.010
57. Croes, K., Colles, A., Koppen, G., Govarts, E., Bruckers, L., Van de Mierop, E., Nelen, V., et al. (2012). Persistent organic pollutants (POPs) in human milk: a biomonitoring study in rural areas of Flanders (Belgium). *Chemosphere*, 89(8), 988–94. doi:10.1016/j.chemosphere.2012.06.058
58. Cunha, S. C., Kalachova, K., Pulkrabova, J., Fernandes, J. O., Oliveira, M. B. P. P., Alves, A., & Hajslova, J. (2010). Polybrominated diphenyl ethers (PBDEs) contents in house and car dust of Portugal by pressurized liquid extraction (PLE) and gas chromatography-mass spectrometry (GC-MS). *Chemosphere*, 78(10), 1263–71. doi:10.1016/j.chemosphere.2009.12.037
59. D'Hollander, W., Roosens, L., Covaci, A., Cornelis, C., Reynders, H., Campenhout, K. Van, Voogt, P. de, et al. (2010). Brominated flame retardants and perfluorinated compounds in indoor dust from homes and offices in Flanders, Belgium. *Chemosphere*, 81(4), 478–87. doi:10.1016/j.chemosphere.2010.07.043
60. Daniels, J. L., Pan, I.-J., Jones, R., Anderson, S., Patterson, D. G., Needham, L. L., & Sjödin, A. (2010). Individual characteristics associated with PBDE levels in U.S. human milk samples. *Environmental health perspectives*, 118(1), 155–60. doi:10.1289/ehp.0900759
61. Darnerud, P. (2003). Toxic effects of brominated flame retardants in man and in wildlife. *Environment International*, 29(6), 841–853. doi:10.1016/S0160-4120(03)00107-7
62. Darnerud, P. O., Aune, M., Larsson, L., Lignell, S., Mutshatshi, T., Okonkwo, J., Botha, B., et al. (2011). Levels of brominated flame retardants and other persistent organic pollutants in breast milk samples from Limpopo Province, South Africa. *The Science of the total environment*, 409(19), 4048–53. doi:10.1016/j.scitotenv.2011.05.054

63. Daubié, S., Bisson, J.-F., Lalonde, R., Schroeder, H., & Rychen, G. (2011). Neurobehavioral and physiological effects of low doses of polybrominated diphenyl ether (PBDE)-99 in male adult rats. *Toxicology letters*, *204*(1), 57–63. doi:10.1016/j.toxlet.2011.04.009
64. De Coster, S., & Van Larebeke, N. (2012). Endocrine-disrupting chemicals: associated disorders and mechanisms of action. *Journal of environmental and public health*, *2012*, 713696. doi:10.1155/2012/713696
65. De Wit, C. A., Björklund, J. A., & Thuresson, K. (2012). Tri-decabrominated diphenyl ethers and hexabromocyclododecane in indoor air and dust from Stockholm microenvironments 2: indoor sources and human exposure. *Environment international*, *39*(1), 141–7. doi:10.1016/j.envint.2011.11.001
66. Devanathan, G., Subramanian, A., Sudaryanto, A., Takahashi, S., Isobe, T., & Tanabe, S. (2012). Brominated flame retardants and polychlorinated biphenyls in human breast milk from several locations in India: potential contaminant sources in a municipal dumping site. *Environment international*, *39*(1), 87–95. doi:10.1016/j.envint.2011.10.005
67. DiGangi, J., Blum, A., Bergman, A., De Wit, C. A., Lucas, D., Mortimer, D., Schecter, A., et al. (2010). San Antonio Statement on brominated and chlorinated flame retardants. *Environmental health perspectives*, *118*(12), A516–8. doi:10.1289/ehp.1003089
68. Dingemans, M. M. L., Van den Berg, M., Bergman, A., & Westerink, R. H. S. (2010). Calcium-related processes involved in the inhibition of depolarization-evoked calcium increase by hydroxylated PBDEs in PC12 cells. *Toxicological sciences*: an official journal of the Society of Toxicology, *114*(2), 302–9. doi:10.1093/toxsci/kfp310
69. Dingemans, M. M. L., Van den Berg, M., & Westerink, R. H. S. (2011). Neurotoxicity of brominated flame retardants: (in)direct effects of parent and hydroxylated polybrominated diphenyl ethers on the (developing) nervous system. *Environmental health perspectives*, *119*(7), 900–7. doi:10.1289/ehp.1003035
70. Dirtu, A. C., Ali, N., Van den Eede, N., Neels, H., & Covaci, A. (2012). Country specific comparison for profile of chlorinated, brominated and phosphate organic contaminants in indoor dust. Case study for Eastern Romania, 2010. *Environment international*, *49*, 1–8. doi:10.1016/j.envint.2012.08.002
71. Dishaw, L. V., Powers, C. M., Ryde, I. T., Roberts, S. C., Seidler, F. J., Slotkin, T. A., & Stapleton, H. M. (2011). Is the PentaBDE Replacement, Tris (1,3-dichloro-2-propyl) Phosphate (TDCPP), a Developmental Neurotoxicant? Studies in PC12 Cells. *Toxicology and Applied Pharmacology*. doi:10.1016/j.taap.2011.01.005
72. Dodson, R. E., Perovich, L. J., Covaci, A., Van den Eede, N., Ionas, A. C., Dirtu, A. C., Brody, J. G., et al. (2012). After the PBDE phase-out: a broad suite of flame retardants in repeat house dust samples from California. *Environmental science & technology*, *46*(24), 13056–66. doi:10.1021/es303879n
73. Domingo, J. L. (2012). Polybrominated diphenyl ethers in food and human dietary exposure: a review of the recent scientific literature. *Food and chemical toxicology*: an international journal published for the British Industrial Biological Research Association, *50*(2), 238–49. doi:10.1016/j.fct.2011.11.004
74. Dreiem, A., Okoniewski, R. J., Brosch, K. O., Miller, V. M., & Seegal, R. F. (2010). Polychlorinated biphenyls and polybrominated diphenyl ethers alter striatal dopamine neurochemistry in synaptosomes from developing rats in an additive manner. *Toxicological sciences*: an official journal of the Society of Toxicology, *118*(1), 150–9. doi:10.1093/toxsci/kfq229
75. Driscoll, L. L., Kaplan, J., Bucuvalas, E., Allen, H., Kraut, J., & Fitzpatrick, J. (n.d.). Acute postnatal exposure to the pentaBDE commercial mixture DE-71 at 5 or 15 mg/kg/day does not produce learning or attention deficits in rats. *Neurotoxicology and teratology*, *34*(1), 20–6. doi:10.1016/j.ntt.2011.10.002
76. Dye, J. A., Venier, M., Zhu, L., Ward, C. R., Hites, R. A., & Birnbaum, L. S. (2007). Elevated PBDE Levels in Pet Cats: Sentinels for Humans? *Environmental Science & Technology*, *41*(18), 6350–6356. doi:10.1021/es0708159
77. Eggesbø, M., Thomsen, C., Jørgensen, J. V., Becher, G., Odland, J. Ø., & Longnecker, M. P. (2011). Associations between brominated flame retardants in human milk and thyroid-stimulating hormone (TSH) in neonates. *Environmental research*, *111*(6), 737–43. doi:10.1016/j.envres.2011.05.004

78. Eriksson, P., Jakobsson, E., & Fredriksson, A. (2001). Brominated flame retardants: a novel class of developmental neurotoxicants in our environment? *Environmental health perspectives*, 109(9), 903–8. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1240439&tool=pmcentrez&rendertype=abstract>
79. Eriksson, P., Viberg, H., Jakobsson, E., Orn, U., & Fredriksson, A. (2002). A brominated flame retardant, 2,2',4,4',5-pentabromodiphenyl ether: uptake, retention, and induction of neurobehavioral alterations in mice during a critical phase of neonatal brain development. *Toxicological sciences*, 67(1), 98–103. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11961221>
80. Eriksson, Per, Fischer, C., & Fredriksson, A. (2006). Polybrominated diphenyl ethers, a group of brominated flame retardants, can interact with polychlorinated biphenyls in enhancing developmental neurobehavioral defects. *Toxicological sciences*, 94(2), 302–9. doi:10.1093/toxsci/kfl109
81. Ernest, S. R., Wade, M. G., Lalancette, C., Ma, Y.-Q., Berger, R. G., Robaire, B., & Hales, B. F. (2012). Effects of chronic exposure to an environmentally relevant mixture of brominated flame retardants on the reproductive and thyroid system in adult male rats. *Toxicological sciences* □: *an official journal of the Society of Toxicology*, 127(2), 496–507. doi:10.1093/toxsci/kfs098
82. Erratico, C. A., Moffatt, S. C., & Bandiera, S. M. (2011). Comparative oxidative metabolism of BDE-47 and BDE-99 by rat hepatic microsomes. *Toxicological sciences* □: *an official journal of the Society of Toxicology*, 123(1), 37–47. doi:10.1093/toxsci/kfr155
83. Erratico, C. A., Szeitz, A., & Bandiera, S. M. (2013). Biotransformation of 2,2',4,4'-Tetrabromodiphenyl Ether (BDE-47) by Human Liver Microsomes: Identification of Cytochrome P450 2B6 as the Major Enzyme Involved. *Chemical research in toxicology*, 26(5), 721–31. doi:10.1021/tx300522u
84. Eskenazi, B., Chevrier, J., Rauch, S. A., Kogut, K., Harley, K. G., Johnson, C., Trujillo, C., et al. (2013). In utero and childhood polybrominated diphenyl ether (PBDE) exposures and neurodevelopment in the CHAMACOS study. *Environmental health perspectives*, 121(2), 257–62. doi:10.1289/ehp.1205597
85. Eskenazi, B., Fenster, L., Castorina, R., Marks, A. R., Sjödin, A., Rosas, L. G., Holland, N., et al. (2011). A comparison of PBDE serum concentrations in Mexican and Mexican-American children living in California. *Environmental health perspectives*, 119(10), 1442–8. doi:10.1289/ehp.1002874
86. Fair, P. A., Stavros, H.-C., Mollenhauer, M. A. M., DeWitt, J. C., Henry, N., Kannan, K., Yun, S. H., et al. (2012). Immune function in female B(6)C(3)F(1) mice is modulated by DE-71, a commercial polybrominated diphenyl ether mixture. *Journal of immunotoxicology*, 9(1), 96–107. doi:10.3109/1547691X.2011.643418
87. Fan, C.-Y., Besas, J., & Kodavanti, P. R. S. (2010). Changes in mitogen-activated protein kinase in cerebellar granule neurons by polybrominated diphenyl ethers and polychlorinated biphenyls. *Toxicology and applied pharmacology*, 245(1), 1–8. doi:10.1016/j.taap.2010.02.008
88. Fang, M., Webster, T. F., Gooden, D., Cooper, E. M., McClean, M. D., Carignan, C., Makey, C., et al. (2013). Investigating a novel flame retardant known as v6: measurements in baby products, house dust, and car dust. *Environmental science & technology*, 47(9), 4449–54. doi:10.1021/es400032v
89. Fängström, B., Athanassiadis, I., Odsjö, T., Norén, K., & Bergman, A. (2008). Temporal trends of polybrominated diphenyl ethers and hexabromocyclododecane in milk from Stockholm mothers, 1980-2004. *Molecular nutrition & food research*, 52(2), 187–93. doi:10.1002/mnfr.200700182
90. Fernandes, A., Dicks, P., Mortimer, D., Gem, M., Smith, F., Driffield, M., White, S., et al. (2008). Brominated and chlorinated dioxins, PCBs and brominated flame retardants in Scottish shellfish: methodology, occurrence and human dietary exposure. *Molecular nutrition & food research*, 52(2), 238–49. doi:10.1002/mnfr.200700135
91. Fischer, D., Hooper, K., Athanasiadou, M., Athanassiadis, I., & Bergman, A. (2006). Children show highest levels of polybrominated diphenyl ethers in a California family of four: a case study. *Environmental health perspectives*, 114(10), 1581–4. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1626410&tool=pmcentrez&rendertype=abstract>

92. Foster, W. G., Gregorovich, S., Morrison, K. M., Atkinson, S. A., Kubwabo, C., Stewart, B., & Teo, K. (2011). Human maternal and umbilical cord blood concentrations of polybrominated diphenyl ethers. *Chemosphere*, *84*(10), 1301–9. doi:10.1016/j.chemosphere.2011.05.028
93. Frederiksen, M., Thomsen, C., Frøshaug, M., Vorkamp, K., Thomsen, M., Becher, G., & Knudsen, L. E. (2010). Polybrominated diphenyl ethers in paired samples of maternal and umbilical cord blood plasma and associations with house dust in a Danish cohort. *International journal of hygiene and environmental health*, *213*(4), 233–42. doi:10.1016/j.ijheh.2010.04.008
94. Frederiksen, M., Vorkamp, K., Mathiesen, L., Mose, T., & Knudsen, L. E. (2010). Placental transfer of the polybrominated diphenyl ethers BDE-47, BDE-99 and BDE-209 in a human placenta perfusion system: an experimental study. *Environmental health: a global access science source*, *9*, 32. doi:10.1186/1476-069X-9-32
95. Frederiksen, M., Vorkamp, K., Thomsen, M., & Knudsen, L. E. (2009). Human internal and external exposure to PBDEs--a review of levels and sources. *International journal of hygiene and environmental health*, *212*(2), 109–34. doi:10.1016/j.ijheh.2008.04.005
96. Fujii, Y., Ito, Y., Harada, K. H., Hitomi, T., Koizumi, A., & Haraguchi, K. (2012). Regional variation and possible sources of brominated contaminants in breast milk from Japan. *Environmental pollution (Barking, Essex) : 1987*, *162*, 269–74. doi:10.1016/j.envpol.2011.11.022
97. Fujimoto, H., Woo, G.-H., Inoue, K., Takahashi, M., Hirose, M., Nishikawa, A., & Shibutani, M. (2011). Impaired oligodendroglial development by decabromodiphenyl ether in rat offspring after maternal exposure from mid-gestation through lactation. *Reproductive toxicology (Elmsford, N.Y.)*, *31*(1), 86–94. doi:10.1016/j.reprotox.2010.09.003
98. Fulong, C. R. P., & Espino, M. P. B. (2013). Decabromodiphenyl ether in indoor dust from different microenvironments in a university in the Philippines. *Chemosphere*, *90*(1), 42–8. doi:10.1016/j.chemosphere.2012.06.072
99. Gascon, M., Vrijheid, M., Martínez, D., Forn, J., Grimalt, J. O., Torrent, M., & Sunyer, J. (2011). Effects of pre and postnatal exposure to low levels of polybromodiphenyl ethers on neurodevelopment and thyroid hormone levels at 4 years of age. *Environment international*, *37*(3), 605–11. doi:10.1016/j.envint.2010.12.005
100. Gaylor, M. O., Harvey, E., & Hale, R. C. (2012). House crickets can accumulate polybrominated diphenyl ethers (PBDEs) directly from polyurethane foam common in consumer products. *Chemosphere*, *86*(5), 500–5. doi:10.1016/j.chemosphere.2011.10.014
101. Gee, J. R., Moser, V. C., McDanie, K. L., & Herr, D. W. (2011). Neurochemical changes following a single dose of polybrominated diphenyl ether 47 in mice. *Drug and chemical toxicology*, *34*(2), 213–9. doi:10.3109/01480545.2010.536768
102. Giordano, G., & Costa, L. G. (2012). Developmental neurotoxicity: some old and new issues. *ISRN toxicology*, *2012*, 814795. doi:10.5402/2012/814795
103. Gold, M. D., Blum, A., & Ames, B. N. (1978). Another flame retardant, tris-(1,3-dichloro-2-propyl)-phosphate, and its expected metabolites are mutagens. *Science*, *200*(4343), 785–7. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/347576>
104. Guo, W., Park, J.-S., Wang, Y., Gardner, S., Baek, C., Petreas, M., & Hooper, K. (2012). High polybrominated diphenyl ether levels in California house cats: house dust a primary source? *Environmental toxicology and chemistry / SETAC*, *31*(2), 301–6. doi:10.1002/etc.1700
105. Hamers, T., Kamstra, J. H., Sonneveld, E., Murk, A. J., Kester, M. H. A., Andersson, P. L., Legler, J., et al. (2006). In vitro profiling of the endocrine-disrupting potency of brominated flame retardants. *Toxicological sciences*, *92*(1), 157–73. doi:10.1093/toxsci/kfj187
106. Han, X., Tang, R., Chen, X., Xu, B., Qin, Y., Wu, W., Hu, Y., et al. (2012). 2,2',4,4'-Tetrabromodiphenyl ether (BDE-47) decreases progesterone synthesis through cAMP-PKA pathway and P450scc downregulation in mouse Leydig tumor cells. *Toxicology*, *302*(1), 44–50. doi:10.1016/j.tox.2012.07.010

107. Hardy, M. L., Banasik, M., & Stedeford, T. (2009). Toxicology and human health assessment of decabromodiphenyl ether. *Critical reviews in toxicology*, *39 Suppl 3*, 1–44. doi:10.1080/10408440902837967
108. Hardy, M. L., Mercieca, M. D., Rodwell, D. E., & Stedeford, T. (2010). Prenatal developmental toxicity of decabromodiphenyl ethane in the rat and rabbit. *Birth defects research. Part B, Developmental and reproductive toxicology*, *89(2)*, 139–46. doi:10.1002/bdrb.20237
109. Harley, K. G., Chevrier, J., Schall, R. A., Sjödin, A., Bradman, A., & Eskenazi, B. (2011). Association of prenatal exposure to polybrominated diphenyl ethers and infant birth weight. *American journal of epidemiology*, *174(8)*, 885–92. doi:10.1093/aje/kwr212
110. Harley, K. G., Marks, A. R., Chevrier, J., Bradman, A., Sjödin, A., & Eskenazi, B. (2010). PBDE concentrations in women's serum and fecundability. *Environmental health perspectives*, *118(5)*, 699–704. doi:10.1289/ehp.0901450
111. Harrad, S., De Wit, C. A., Abdallah, M. A.-E., Bergh, C., Björklund, J. A., Covaci, A., Darnerud, P. O., et al. (2010). Indoor contamination with hexabromocyclododecanes, polybrominated diphenyl ethers, and perfluoroalkyl compounds: an important exposure pathway for people? *Environmental science & technology*, *44(9)*, 3221–31. doi:10.1021/es903476t
112. Harrad, S., Goosey, E., Desborough, J., Abdallah, M. A.-E., Roosens, L., & Covaci, A. (2010). Dust from U.K. primary school classrooms and daycare centers: the significance of dust as a pathway of exposure of young U.K. children to brominated flame retardants and polychlorinated biphenyls. *Environmental science & technology*, *44(11)*, 4198–202. doi:10.1021/es100750s
113. Harrad, S., Hazrati, S., & Ibarra, C. (2006). Concentrations of Polychlorinated Biphenyls in Indoor Air and Polybrominated Diphenyl Ethers in Indoor Air and Dust in Birmingham, United Kingdom: Implications for Human Exposure. *Environmental Science & Technology*, *40(15)*, 4633–4638. doi:10.1021/es0609147
114. Hashimoto, S., Yoshimura, H., Okada, K., Uramaru, N., Sugihara, K., Kitamura, S., & Imaoka, S. (2012). Effects of polybrominated diphenyl ethers (PBDEs) and their derivatives on protein disulfide isomerase activity and growth hormone release of GH3 cells. *Chemical research in toxicology*, *25(3)*, 656–63. doi:10.1021/tx200374s
115. He, P., Wang, A., Niu, Q., Guo, L., Xia, T., & Chen, X. (2011). Toxic effect of PBDE-47 on thyroid development, learning, and memory, and the interaction between PBDE-47 and PCB153 that enhances toxicity in rats. *Toxicology and industrial health*, *27(3)*, 279–88. doi:10.1177/07482337110387002
116. He, P., Wang, A., Xia, T., Gao, P., Niu, Q., Guo, L., & Chen, X. (2009). Mechanisms underlying the developmental neurotoxic effect of PBDE-47 and the enhanced toxicity associated with its combination with PCB153 in rats. *Neurotoxicology*, *30(6)*, 1088–95. doi:10.1016/j.neuro.2009.06.005
117. Helleday, T., Tuominen, K. L., Bergman, A., & Jessen, D. (1999). Brominated flame retardants induce intragenic recombination in mammalian cells. *Mutation research*, *439(2)*, 137–47. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10023042>
118. Herbstman, J. B., Sjödin, A., Kurzon, M., Lederman, S. A., Jones, R. S., Rauh, V., Needham, L. L., et al. (2010). Prenatal exposure to PBDEs and neurodevelopment. *Environmental health perspectives*, *118(5)*, 712–9. doi:10.1289/ehp.0901340
119. Heredia, L., Torrente, M., Colomina, M. T., & Domingo, J. L. (2012). Behavioral effects of oral subacute exposure to BDE-209 in young adult mice: a preliminary study. *Food and chemical toxicology*: an international journal published for the British Industrial Biological Research Association, *50(3-4)*, 707–12. doi:10.1016/j.fct.2011.12.002
120. Hertz-Picciotto, I., Bergman, A., Fängström, B., Rose, M., Krakowiak, P., Pessah, I., Hansen, R., et al. (2011). Polybrominated diphenyl ethers in relation to autism and developmental delay: a case-control study. *Environmental health*: a global access science source, *10(1)*, 1. doi:10.1186/1476-069X-10-1
121. Hirai, T., Fujimine, Y., Watanabe, S., & Nakano, T. (2012). Distribution of polybrominated diphenyl ethers in Japanese autopsy tissue and body fluid samples. *Environmental science and pollution research international*, *19(8)*, 3538–46. doi:10.1007/s11356-012-0915-z

122. Hites, R. A. (2004). Polybrominated diphenyl ethers in the environment and in people: a meta-analysis of concentrations. *Environmental science & technology*, 38(4), 945–56. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14998004>
123. Ho, K.-L., Murphy, M. B., Wan, Y., Fong, B. M.-W., Tam, S., Giesy, J. P., Leung, K. S.-Y., et al. (2012). Synthesis and characterization of bromophenol glucuronide and sulfate conjugates for their direct LC-MS/MS quantification in human urine as potential exposure markers for polybrominated diphenyl ethers. *Analytical chemistry*, 84(22), 9881–8. doi:10.1021/ac302161t
124. Hoffman, K., Adgent, M., Goldman, B. D., Sjödin, A., & Daniels, J. L. (2012). Lactational exposure to polybrominated diphenyl ethers and its relation to social and emotional development among toddlers. *Environmental health perspectives*, 120(10), 1438–42. doi:10.1289/ehp.1205100
125. Hoppe, A. A., & Carey, G. B. (2007). Polybrominated diphenyl ethers as endocrine disruptors of adipocyte metabolism. *Obesity*, 15(12), 2942–50. doi:10.1038/oby.2007.351
126. Horton, M. K., Bousleiman, S., Jones, R., Sjödin, A., Liu, X., Whyatt, R., Wapner, R., et al. (2013). Predictors of serum concentrations of polybrominated flame retardants among healthy pregnant women in an urban environment: a cross-sectional study. *Environmental health : a global access science source*, 12(1), 23. doi:10.1186/1476-069X-12-23
127. Huang, S. C., Giordano, G., & Costa, L. G. (2010). Comparative cytotoxicity and intracellular accumulation of five polybrominated diphenyl ether congeners in mouse cerebellar granule neurons. *Toxicological sciences : an official journal of the Society of Toxicology*, 114(1), 124–32. doi:10.1093/toxsci/kfp296
128. Hudec, T., Thean, J., Kuehl, D., & Dougherty, R. (1981). Tris(dichloropropyl)phosphate, a mutagenic flame retardant: frequent cocurrence in human seminal plasma. *Science*, 211(4485), 951–952. doi:10.1126/science.7466368
129. Ibhazehiebo, K., Iwasaki, T., Kimura-Kuroda, J., Miyazaki, W., Shimokawa, N., & Koibuchi, N. (2011). Disruption of thyroid hormone receptor-mediated transcription and thyroid hormone-induced Purkinje cell dendrite arborization by polybrominated diphenyl ethers. *Environmental health perspectives*, 119(2), 168–75. doi:10.1289/ehp.1002065
130. Imm, P., Knobeloch, L., Buelow, C., & Anderson, H. A. (2009). Household exposures to polybrominated diphenyl ethers (PBDEs) in a Wisconsin Cohort. *Environmental health perspectives*, 117(12), 1890–5. doi:10.1289/ehp.0900839
131. Ji, K., Choi, K., Giesy, J. P., Musarrat, J., & Takeda, S. (2011). Genotoxicity of several polybrominated diphenyl ethers (PBDEs) and hydroxylated PBDEs, and their mechanisms of toxicity. *Environmental science & technology*, 45(11), 5003–8. doi:10.1021/es104344e
132. Jin, J., Wang, Y., Yang, C., Hu, J., Liu, W., Cui, J., & Tang, X. (2009). Polybrominated diphenyl ethers in the serum and breast milk of the resident population from production area, China. *Environment international*, 35(7), 1048–52. doi:10.1016/j.envint.2009.05.006
133. Johansson, N., Viberg, H., Fredriksson, A., & Eriksson, P. (2008). Neonatal exposure to deca-brominated diphenyl ether (PBDE 209) causes dose-response changes in spontaneous behaviour and cholinergic susceptibility in adult mice. *Neurotoxicology*, 29(6), 911–9. doi:10.1016/j.neuro.2008.09.008
134. Johnson, P. I., Stapleton, H. M., Mukherjee, B., Hauser, R., & Meeker, J. D. (2013). Associations between brominated flame retardants in house dust and hormone levels in men. *The Science of the total environment*, 445–446, 177–84. doi:10.1016/j.scitotenv.2012.12.017
135. Johnson-Restrepo, B., Addink, R., Wong, C., Arcaro, K., & Kannan, K. (2007). Polybrominated diphenyl ethers and organochlorine pesticides in human breast milk from Massachusetts, USA. *Journal of environmental monitoring : JEM*, 9(11), 1205–12. doi:10.1039/b711409p
136. Johnson-Restrepo, B., & Kannan, K. (2009). An assessment of sources and pathways of human exposure to polybrominated diphenyl ethers in the United States. *Chemosphere*, 76(4), 542–8. doi:10.1016/j.chemosphere.2009.02.068

137. Johnson-Restrepo, B., Kannan, K., Rapaport, D. P., & Rodan, B. D. (2005). Polybrominated diphenyl ethers and polychlorinated biphenyls in human adipose tissue from New York. *Environmental science & technology*, 39(14), 5177–82. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16082945>
138. Jones-Otazo, H. A., Clarke, J. P., Diamond, M. L., Archbold, J. A., Ferguson, G., Harner, T., Richardson, G. M., et al. (2005). Is House Dust the Missing Exposure Pathway for PBDEs? An Analysis of the Urban Fate and Human Exposure to PBDEs. *Environmental Science & Technology*, 39(14), 5121–5130. doi:10.1021/es048267b
139. Jonsson, O. B., Dyremark, E., & Nilsson, U. L. (2001). Development of a microporous membrane liquid-liquid extractor for organophosphate esters in human blood plasma: identification of triphenyl phosphate and octyl diphenyl phosphate in donor plasma. *Journal of chromatography. B, Biomedical sciences and applications*, 755(1-2), 157–64. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11393700>
140. Jugan, M.-L., Levi, Y., & Blondeau, J.-P. (2010). Endocrine disruptors and thyroid hormone physiology. *Biochemical pharmacology*, 79(7), 939–47. doi:10.1016/j.bcp.2009.11.006
141. Jurewicz, J., Polańska, K., & Hanke, W. (2013). Exposure to widespread environmental toxicants and children's cognitive development and behavioral problems. *International journal of occupational medicine and environmental health*. doi:10.2478/s13382-013-0099-x
142. Kakimoto, K., Nagayoshi, H., Yoshida, J., Akutsu, K., Konishi, Y., Toriba, A., & Hayakawa, K. (2012). Detection of Dechlorane Plus and brominated flame retardants in marketed fish in Japan. *Chemosphere*, 89(4), 416–9. doi:10.1016/j.chemosphere.2012.05.072
143. Kalachova, K., Hradkova, P., Lankova, D., Hajslova, J., & Pulkrabova, J. (2012). Occurrence of brominated flame retardants in household and car dust from the Czech Republic. *The Science of the total environment*, 441, 182–93. doi:10.1016/j.scitotenv.2012.09.061
144. Kalantzi, O I, Brown, F. R., Caleffi, M., Goth-Goldstein, R., & Petreas, M. (2009). Polybrominated diphenyl ethers and polychlorinated biphenyls in human breast adipose samples from Brazil. *Environment international*, 35(1), 113–7. doi:10.1016/j.envint.2008.09.008
145. Kalantzi, Olga I, Geens, T., Covaci, A., & Siskos, P. A. (2011). Distribution of polybrominated diphenyl ethers (PBDEs) and other persistent organic pollutants in human serum from Greece. *Environment international*, 37(2), 349–53. doi:10.1016/j.envint.2010.10.005
146. Kanazawa, A., Saito, I., Araki, A., Takeda, M., Ma, M., Saijo, Y., & Kishi, R. (2010). Association between indoor exposure to semi-volatile organic compounds and building-related symptoms among the occupants of residential dwellings. *Indoor air*, 20(1), 72–84. doi:10.1111/j.1600-0668.2009.00629.x
147. Karpeta, A., & Gregoraszczyk, E. (2010). Mixture of dominant PBDE congeners (BDE-47, -99, -100 and -209) at levels noted in human blood dramatically enhances progesterone secretion by ovarian follicles. *Endocrine regulations*, 44(2), 49–55. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20429633>
148. Kawashiro, Y., Fukata, H., Omori-Inoue, M., Kubonoya, K., Jotaki, T., Takigami, H., Sakai, S., et al. (2008). Perinatal exposure to brominated flame retardants and polychlorinated biphenyls in Japan. *Endocrine journal*, 55(6), 1071–84. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18719292>
149. Kiciński, M., Viaene, M. K., Den Hond, E., Schoeters, G., Covaci, A., Dirtu, A. C., Nelen, V., et al. (2012). Neurobehavioral function and low-level exposure to brominated flame retardants in adolescents: a cross-sectional study. *Environmental health : a global access science source*, 11, 86. doi:10.1186/1476-069X-11-86
150. Kim, J., Kang, J.-H., Park, H., Baek, S.-Y., Kim, Y.-H., & Chang, Y.-S. (2012). Assessment of polybrominated diphenyl ethers (PBDEs) in serum from the Korean general population. *Environmental pollution (Barking, Essex) : 1987*, 164, 46–52. doi:10.1016/j.envpol.2012.01.016
151. Kim, J.-W., Isobe, T., Sudaryanto, A., Malarvannan, G., Chang, K.-H., Muto, M., Prudente, M., et al. (2013). Organophosphorus flame retardants in house dust from the Philippines: occurrence and assessment of human exposure. *Environmental science and pollution research international*, 20(2), 812–22. doi:10.1007/s11356-012-1237-x

152. Kim, T. H., Bang, D. Y., Lim, H. J., Won, A. J., Ahn, M. Y., Patra, N., Chung, K. K., et al. (2012). Comparisons of polybrominated diphenyl ethers levels in paired South Korean cord blood, maternal blood, and breast milk samples. *Chemosphere*, *87*(1), 97–104. doi:10.1016/j.chemosphere.2011.11.074
153. Koenig, C. M., Lango, J., Pessah, I. N., & Berman, R. F. (n.d.). Maternal transfer of BDE-47 to offspring and neurobehavioral development in C57BL/6J mice. *Neurotoxicology and teratology*, *34*(6), 571–80. doi:10.1016/j.ntt.2012.09.005
154. Koh, T.-W., Chih-Cheng Chen, S., Chang-Chien, G.-P., Lin, D.-Y., Chen, F.-A., & Chao, H.-R. (2010). Breast-milk levels of polybrominated diphenyl ether flame retardants in relation to women's age and pre-pregnant body mass index. *International journal of hygiene and environmental health*, *213*(1), 59–65. doi:10.1016/j.ijheh.2009.09.004
155. Kotz, A., Malisch, R., Kypke, K., & Oehme, M. (2005). PBDE, PBDD/F and mixed chlorinated-brominated PXDD/F in pooled human milk samples from different countries. *Organohalogen Compounds*, *67*, 1540–1544. Retrieved from <http://www.dioxin20xx.org/pdfs/2005/05-267.pdf>
156. Kuriyama, S. N., Wanner, A., Fidalgo-Neto, A. A., Talsness, C. E., Koerner, W., & Chahoud, I. (2007). Developmental exposure to low-dose PBDE-99: tissue distribution and thyroid hormone levels. *Toxicology*, *242*(1-3), 80–90. doi:10.1016/j.tox.2007.09.011
157. Lai, Y., Lu, M., Gao, X., Wu, H., & Cai, Z. (2011). New evidence for toxicity of polybrominated diphenyl ethers: DNA adduct formation from quinone metabolites. *Environmental science & technology*, *45*(24), 10720–7. doi:10.1021/es203068f
158. Legler, J. (2008). New insights into the endocrine disrupting effects of brominated flame retardants. *Chemosphere*, *73*(2), 216–22. doi:10.1016/j.chemosphere.2008.04.081
159. Legler, J., Hamers, T., Van Eck van der Sluijs-van de Bor, M., Schoeters, G., Van der Ven, L., Eggesbo, M., Koppe, J., et al. (2011). The OBELIX project: early life exposure to endocrine disruptors and obesity. *The American journal of clinical nutrition*, *94*(6 Suppl), 1933S–1938S. doi:10.3945/ajcn.110.001669
160. Leijds, M. M., Ten Tusscher, G. W., Olie, K., Van Teunenbroek, T., Van Aalderen, W. M. C., De Voogt, P., Vulmsma, T., et al. (2012). Thyroid hormone metabolism and environmental chemical exposure. *Environmental health: a global access science source*, *11 Suppl 1*, S10. doi:10.1186/1476-069X-11-S1-S10
161. Li, F., Xie, Q., Li, X., Li, N., Chi, P., Chen, J., Wang, Z., et al. (2010). Hormone activity of hydroxylated polybrominated diphenyl ethers on human thyroid receptor-beta: in vitro and in silico investigations. *Environmental health perspectives*, *118*(5), 602–6. doi:10.1289/ehp.0901457
162. Li, T., Wang, W., Pan, Y.-W., Xu, L., & Xia, Z. (2013). A Hydroxylated Metabolite of Flame-Retardant PBDE-47 Decreases the Survival, Proliferation, and Neuronal Differentiation of Primary Cultured Adult Neural Stem Cells and Interferes with Signaling of ERK5 MAP Kinase and Neurotrophin 3. *Toxicological sciences: an official journal of the Society of Toxicology*, kft083–. doi:10.1093/toxsci/kft083
163. Liang, S.-X., Gao, H.-X., Zhao, Y.-Y., Ma, X.-M., & Sun, H.-W. (2010). Effects of repeated exposure to decabrominated diphenyl ether (BDE-209) on mice nervous system and its self repair. *Environmental toxicology and pharmacology*, *29*(3), 297–301. doi:10.1016/j.etap.2010.02.005
164. Liberda, E. N., Wainman, B. C., Leblanc, A., Dumas, P., Martin, I., & Tsuji, L. J. S. (2011). Dietary exposure of PBDEs resulting from a subsistence diet in three First Nation communities in the James Bay Region of Canada. *Environment international*. doi:10.1016/j.envint.2010.12.008
165. Lilienthal, H., Hack, A., Roth-Härer, A., Grande, S. W., & Talsness, C. E. (2006). Effects of developmental exposure to 2,2',4,4',5-pentabromodiphenyl ether (PBDE-99) on sex steroids, sexual development, and sexually dimorphic behavior in rats. *Environmental health perspectives*, *114*(2), 194–201. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1367831&tool=pmcentrez&rendertype=abstract>
166. Lim, J.-S., Lee, D.-H., & Jacobs, D. R. (2008). Association of brominated flame retardants with diabetes and metabolic syndrome in the U.S. population, 2003–2004. *Diabetes care*, *31*(9), 1802–7. doi:10.2337/dc08-0850
167. Lin, K. (2008). Joint acute toxicity of tributyl phosphate and triphenyl phosphate to *Daphnia magna*. *Environmental Chemistry Letters*, *7*(4), 309–312. doi:10.1007/s10311-008-0170-1

168. Link, B., Gabrio, T., Mann, V., Schilling, B., Maisner, V., König, M., Flicker-Klein, A., et al. (2012). Polybrominated diphenyl ethers (PBDE) in blood of children in Baden-Württemberg between 2002/03 and 2008/09. *International journal of hygiene and environmental health*, 215(2), 224–8. doi:10.1016/j.ijheh.2011.10.018
169. Lorber, M. (2008). Exposure of Americans to polybrominated diphenyl ethers. *Journal of exposure science & environmental epidemiology*, 18(1), 2–19. doi:10.1038/sj.jes.7500572
170. Lunder, S., Hovander, L., Athanassiadis, I., & Bergman, A. (2010a). Significantly higher polybrominated diphenyl ether levels in young U.S. children than in their mothers. *Environmental science & technology*, 44(13), 5256–62. doi:10.1021/es1009357
171. Lunder, S., Hovander, L., Athanassiadis, I., & Bergman, A. (2010b). Significantly higher polybrominated diphenyl ether levels in young U.S. children than in their mothers. *Environmental science & technology*, 44(13), 5256–62. doi:10.1021/es1009357
172. Lundgren, M., Darnerud, P. O., Molin, Y., Lilienthal, H., Blomberg, J., & Ilbäck, N.-G. (2007). Viral infection and PBDE exposure interact on CYP gene expression and enzyme activities in the mouse liver. *Toxicology*, 242(1–3), 100–8. doi:10.1016/j.tox.2007.09.014
173. Main, K. M., Kiviranta, H., Virtanen, H. E., Sundqvist, E., Tuomisto, J. T., Tuomisto, J., Vartiainen, T., et al. (2007). Flame retardants in placenta and breast milk and cryptorchidism in newborn boys. *Environmental health perspectives*, 115(10), 1519–26. doi:10.1289/ehp.9924
174. Malarvannan, G., Isobe, T., Covaci, A., Prudente, M., & Tanabe, S. (2013). Accumulation of brominated flame retardants and polychlorinated biphenyls in human breast milk and scalp hair from the Philippines: levels, distribution and profiles. *The Science of the total environment*, 442, 366–79. doi:10.1016/j.scitotenv.2012.10.005
175. Mariussen, E., & Fonnum, F. (2003). The effect of brominated flame retardants on neurotransmitter uptake into rat brain synaptosomes and vesicles. *Neurochemistry International*, 43(4–5), 533–542. doi:10.1016/S0197-0186(03)00044-5
176. McDonald, T. A. (2005). Polybrominated diphenylether levels among United States residents: daily intake and risk of harm to the developing brain and reproductive organs. *Integrated environmental assessment and management*, 1(4), 343–54. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16639901>
177. Meeker, J. D., Cooper, E. M., Stapleton, H. M., & Hauser, R. (2013). Urinary metabolites of organophosphate flame retardants: temporal variability and correlations with house dust concentrations. *Environmental health perspectives*, 121(5), 580–5. doi:10.1289/ehp.1205907
178. Meeker, J. D., Johnson, P. I., Camann, D., & Hauser, R. (2009). Polybrominated diphenyl ether (PBDE) concentrations in house dust are related to hormone levels in men. *The Science of the total environment*, 407(10), 3425–9. doi:10.1016/j.scitotenv.2009.01.030
179. Meeker, J. D., & Stapleton, H. M. (2010). House dust concentrations of organophosphate flame retardants in relation to hormone levels and semen quality parameters. *Environmental health perspectives*, 118(3), 318–23. doi:10.1289/ehp.0901332
180. Meerts, I. A., Van Zanden, J. J., Luijckx, E. A., Van Leeuwen-Bol, I., Marsh, G., Jakobsson, E., Bergman, A., et al. (2000). Potent competitive interactions of some brominated flame retardants and related compounds with human transthyretin in vitro. *Toxicological sciences*, 56(1), 95–104. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10869457>
181. Mensching, D. A., Slater, M., Scott, J. W., Ferguson, D. C., & Beasley, V. R. (2012). The feline thyroid gland: a model for endocrine disruption by polybrominated diphenyl ethers (PBDEs)? *Journal of toxicology and environmental health. Part A*, 75(4), 201–12. doi:10.1080/15287394.2012.652054
182. Messer, A. (2010). Mini-review: polybrominated diphenyl ether (PBDE) flame retardants as potential autism risk factors. *Physiology & behavior*, 100(3), 245–9. doi:10.1016/j.physbeh.2010.01.011

183. Miller, M. F., Chernyak, S. M., Domino, S. E., Batterman, S. A., & Loch-Caruso, R. (2012). Concentrations and speciation of polybrominated diphenyl ethers in human amniotic fluid. *The Science of the total environment*, 417-418, 294–8. doi:10.1016/j.scitotenv.2011.11.088
184. Miyaso, H., Nakamura, N., Matsuno, Y., Kawashiro, Y., Komiyama, M., & Mori, C. (2012). Postnatal exposure to low-dose decabromodiphenyl ether adversely affects mouse testes by increasing tyrosine phosphorylation level of cortactin. *The Journal of toxicological sciences*, 37(5), 987–99. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/23038006>
185. Napoli, E., Hung, C., Wong, S., & Giulivi, C. (2013). Toxicity of the flame-retardant BDE-49 on brain mitochondria and neuronal progenitor striatal cells enhanced by a PTEN-deficient background. *Toxicological sciences*: an official journal of the Society of Toxicology, 132(1), 196–210. doi:10.1093/toxsci/kfs339
186. Nash, J. T., Szabo, D. T., & Carey, G. B. (2013). Polybrominated diphenyl ethers alter hepatic phosphoenolpyruvate carboxykinase enzyme kinetics in male Wistar rats: implications for lipid and glucose metabolism. *Journal of toxicology and environmental health. Part A*, 76(2), 142–56. doi:10.1080/15287394.2012.738457
187. Norrgran, J., Jones, B., Lindquist, N.-G., & Bergman, A. (2012). Decabromobiphenyl, polybrominated diphenyl ethers, and brominated phenolic compounds in serum of cats diagnosed with the endocrine disease feline hyperthyroidism. *Archives of environmental contamination and toxicology*, 63(1), 161–8. doi:10.1007/s00244-012-9750-y
188. Öberg, M., Westerholm, E., Fattore, E., Stern, N., Hanberg, A., Haglund, P., Wiberg, K., et al. (2010). Toxicity of Bromkal 70-5DE, a technical mixture of polybrominated diphenyl ethers, following 28 d of oral exposure in rats and impact of analysed impurities. *Chemosphere*, 80(2), 137–143. doi:10.1016/j.chemosphere.2010.04.006
189. Orta-Garcia, S. T., León-Moreno, L. C., González-Vega, C., Dominguez-Cortinas, G., Espinosa-Reyes, G., & Pérez-Maldonado, I. N. (2012). Assessment of the levels of polybrominated diphenyl ethers in blood samples from Guadalajara, Jalisco, Mexico. *Bulletin of environmental contamination and toxicology*, 89(4), 925–9. doi:10.1007/s00128-012-0727-3
190. Pacyniak, E., Roth, M., Hagenbuch, B., & Guo, G. L. (2010). Mechanism of polybrominated diphenyl ether uptake into the liver: PBDE congeners are substrates of human hepatic OATP transporters. *Toxicological sciences*: an official journal of the Society of Toxicology, 115(2), 344–53. doi:10.1093/toxsci/kfq059
191. Patisaul, H. B., Roberts, S. C., Mabrey, N., McCaffrey, K. A., Gear, R. B., Braun, J., Belcher, S. M., et al. (2013). Accumulation and endocrine disrupting effects of the flame retardant mixture Firemaster® 550 in rats: an exploratory assessment. *Journal of biochemical and molecular toxicology*, 27(2), 124–36. doi:10.1002/jbt.21439
192. Patrick, L. (2009). Thyroid disruption: mechanism and clinical implications in human health. *Alternative medicine review*: a journal of clinical therapeutic, 14(4), 326–46. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20030460>
193. Pellacani, C., Buschini, A., Galati, S., Mussi, F., Franzoni, S., & Costa, L. G. (n.d.). Evaluation of DNA damage induced by 2 polybrominated diphenyl ether flame retardants (BDE-47 and BDE-209) in SK-N-MC cells. *International journal of toxicology*, 31(4), 372–9. doi:10.1177/1091581812449663
194. Pereira, L. C., De Souza, A. O., & Dorta, D. J. (2013). Polybrominated Diphenyl Ether Congener (BDE-100) Induces Mitochondrial Impairment. *Basic & clinical pharmacology & toxicology*, 112(6), 418–24. doi:10.1111/bcpt.12046
195. Pérez-Maldonado, I. N., Ramírez-Jiménez, M. D. R., Martínez-Arévalo, L. P., López-Guzmán, O. D., Athanasiadou, M., Bergman, A., Yarto-Ramírez, M., et al. (2009). Exposure assessment of polybrominated diphenyl ethers (PBDEs) in Mexican children. *Chemosphere*, 75(9), 1215–20. doi:10.1016/j.chemosphere.2009.01.083
196. Petreas, M., Nelson, D., Brown, F. R., Goldberg, D., Hurley, S., & Reynolds, P. (2011). High concentrations of polybrominated diphenylethers (PBDEs) in breast adipose tissue of California women. *Environment international*, 37(1), 190–7. doi:10.1016/j.envint.2010.09.001

197. Petreas, M., She, J., Brown, F. R., Winkler, J., Windham, G., Rogers, E., Zhao, G., et al. (2003). High body burdens of 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) in California women. *Environmental health perspectives*, *111*(9), 1175–9. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1241571&tool=pmcentrez&rendertype=abstract>
198. Quirós-Alcalá, L., Bradman, A., Nishioka, M., Harnly, M. E., Hubbard, A., McKone, T. E., & Eskenazi, B. (2011). Concentrations and loadings of polybrominated diphenyl ethers in dust from low-income households in California. *Environment international*. doi:10.1016/j.envint.2010.12.003
199. Rahman, F., Langford, K. H., Scrimshaw, M. D., & Lester, J. N. (2001). Polybrominated diphenyl ether (PBDE) flame retardants. *The Science of the total environment*, *275*(1-3), 1–17. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11482396>
200. Ravnum, S., Zimmer, K. E., Keune, H., Gutleb, A. C., Murk, A. J., Koppe, J. G., Magnanti, B., et al. (2012). Policy relevant results from an expert elicitation on the human health risks of decabromodiphenyl ether (decaBDE) and hexabromocyclododecane (HBCD). *Environmental health perspectives: a global access science source*, *11 Suppl 1*, S7. doi:10.1186/1476-069X-11-S1-S7
201. Reistad, T., Fonnum, F., & Mariussen, E. (2006). Neurotoxicity of the pentabrominated diphenyl ether mixture, DE-71, and hexabromocyclododecane (HBCD) in rat cerebellar granule cells in vitro. *Archives of toxicology*, *80*(11), 785–96. doi:10.1007/s00204-006-0099-8
202. Ren, X., Han, H. J., Lee, Y. J., Lee, S. H., Ng, H. Y., Chae, K.-J., & Kim, I. S. (2012). Proapoptotic effect of a micropollutant (tris-(2-chloroethyl)-phosphate) at environmental level in primary cultured renal proximal tubule cells. *Journal of water and health*, *10*(4), 522–30. doi:10.2166/wh.2012.180
203. Rice, D. C., Reeve, E. A., Herlihy, A., Zoeller, R. T., Thompson, W. D., & Markowski, V. P. (n.d.). Developmental delays and locomotor activity in the C57BL6/J mouse following neonatal exposure to the fully-brominated PBDE, decabromodiphenyl ether. *Neurotoxicology and teratology*, *29*(4), 511–20. doi:10.1016/j.ntt.2007.03.061
204. Rice, D. C., Thompson, W. D., Reeve, E. A., Onos, K. D., Assadollahzadeh, M., & Markowski, V. P. (2009). Behavioral changes in aging but not young mice after neonatal exposure to the polybrominated flame retardant decaBDE. *Environmental health perspectives*, *117*(12), 1903–11. doi:10.1289/ehp.11814
205. Roberts, S. C., Macaulay, L. J., & Stapleton, H. M. (2012). In vitro metabolism of the brominated flame retardants 2-ethylhexyl-2,3,4,5-tetrabromobenzoate (TBB) and bis(2-ethylhexyl) 2,3,4,5-tetrabromophthalate (TBPH) in human and rat tissues. *Chemical research in toxicology*, *25*(7), 1435–41. doi:10.1021/tx300086x
206. Roosens, L., Cornelis, C., D'Hollander, W., Bervoets, L., Reynders, H., Van Campenhout, K., Van Den Heuvel, R., et al. (2010). Exposure of the Flemish population to brominated flame retardants: model and risk assessment. *Environment international*, *36*(4), 368–76. doi:10.1016/j.envint.2010.02.005
207. Roosens, L., D'Hollander, W., Bervoets, L., Reynders, H., Van Campenhout, K., Cornelis, C., Van Den Heuvel, R., et al. (2010). Brominated flame retardants and perfluorinated chemicals, two groups of persistent contaminants in Belgian human blood and milk. *Environmental pollution (Barking, Essex): 1987*, *158*(8), 2546–52. doi:10.1016/j.envpol.2010.05.022
208. Rose, M., Bennett, D. H., Bergman, A., Fängström, B., Pessah, I. N., & Hertz-Picciotto, I. (2010). PBDEs in 2–5 year-old children from California and associations with diet and indoor environment. *Environmental science & technology*, *44*(7), 2648–53. doi:10.1021/es903240g
209. Roze, E., Meijer, L., Bakker, A., Van Braeckel, K. N. J. A., Sauer, P. J. J., & Bos, A. F. (2009). Prenatal exposure to organohalogenes, including brominated flame retardants, influences motor, cognitive, and behavioral performance at school age. *Environmental health perspectives*, *117*(12), 1953–8. doi:10.1289/ehp.0901015
210. Rudel, R. A., Dodson, R. E., Perovich, L. J., Morello-Frosch, R., Camann, D. E., Zuniga, M. M., Yau, A. Y., et al. (2010). Semivolatile endocrine-disrupting compounds in paired indoor and outdoor air in two northern California communities. *Environmental science & technology*, *44*(17), 6583–90. doi:10.1021/es100159c

211. Schecter, A., Colacino, J., Sjödin, A., Needham, L., & Birnbaum, L. (2010). Partitioning of polybrominated diphenyl ethers (PBDEs) in serum and milk from the same mothers. *Chemosphere*, 78(10), 1279–84. doi:10.1016/j.chemosphere.2009.12.016
212. Schecter, A., Haffner, D., Colacino, J., Patel, K., Päpke, O., Opel, M., & Birnbaum, L. (2010). Polybrominated diphenyl ethers (PBDEs) and hexabromocyclodecane (HBCD) in composite U.S. food samples. *Environmental health perspectives*, 118(3), 357–62. doi:10.1289/ehp.0901345
213. Schecter, A., Harris, T. R., Shah, N., Musumba, A., & Päpke, O. (2008). Brominated flame retardants in US food. *Molecular nutrition & food research*, 52(2), 266–72. doi:10.1002/mnfr.200700166
214. Schreiber, T., Gassmann, K., Götz, C., Hübenthal, U., Moors, M., Krause, G., Merk, H. F., et al. (2010). Polybrominated diphenyl ethers induce developmental neurotoxicity in a human in vitro model: evidence for endocrine disruption. *Environmental health perspectives*, 118(4), 572–8. doi:10.1289/ehp.0901435
215. Shaw, S. D., Blum, A., Weber, R., Kannan, K., Rich, D., Lucas, D., Koshland, C. P., et al. (2010). Halogenated flame retardants: do the fire safety benefits justify the risks? *Reviews on environmental health*, 25(4), 261–305. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21268442>
216. She, J., Petreas, M., Winkler, J., Visita, P., McKinney, M., & Kopec, D. (2002). PBDEs in the San Francisco Bay Area: measurements in harbor seal blubber and human breast adipose tissue. *Chemosphere*, 46(5), 697–707. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11999793>
217. Shi, Z., Jiao, Y., Hu, Y., Sun, Z., Zhou, X., Feng, J., Li, J., et al. (2013). Levels of tetrabromobisphenol A, hexabromocyclododecanes and polybrominated diphenyl ethers in human milk from the general population in Beijing, China. *The Science of the total environment*, 452–453, 10–8. doi:10.1016/j.scitotenv.2013.02.038
218. Shoeib, M., Harner, T., Webster, G. M., Sverko, E., & Cheng, Y. (2012). Legacy and current-use flame retardants in house dust from Vancouver, Canada. *Environmental pollution (Barking, Essex 1987)*, 169, 175–82. doi:10.1016/j.envpol.2012.01.043
219. Siddique, S., Xian, Q., Abdelouahab, N., Takser, L., Phillips, S. P., Feng, Y.-L., Wang, B., et al. (2012). Levels of dechlorane plus and polybrominated diphenylethers in human milk in two Canadian cities. *Environment international*, 39(1), 50–5. doi:10.1016/j.envint.2011.09.010
220. Sjödin, A., Patterson, D. G., & Bergman, A. (2001). Brominated flame retardants in serum from U.S. blood donors. *Environmental science & technology*, 35(19), 3830–3. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11642440>
221. Sjödin, A., Päpke, O., McGahee, E., Focant, J.-F., Jones, R. S., Pless-Mulloli, T., Toms, L.-M. L., et al. (2008). Concentration of polybrominated diphenyl ethers (PBDEs) in household dust from various countries. *Chemosphere*, 73(1 Suppl), S131–6. doi:10.1016/j.chemosphere.2007.08.075
222. Small, C. M., Murray, D., Terrell, M. L., & Marcus, M. (2011). Reproductive outcomes among women exposed to a brominated flame retardant in utero. *Archives of environmental & occupational health*, 66(4), 201–8. doi:10.1080/19338244.2010.539640
223. Souza, A. O., Pereira, L. C., Oliveira, D. P., & Dorta, D. J. (2013). BDE-99 congener induces cell death by apoptosis of human hepatoblastoma cell line - HepG2. *Toxicology in vitro: an international journal published in association with BIBRA*, 27(2), 580–7. doi:10.1016/j.tiv.2012.09.022
224. Springer, C., Dere, E., Hall, S. J., McDonnell, E. V., Roberts, S. C., Butt, C. M., Stapleton, H. M., et al. (2012). Rodent thyroid, liver, and fetal testis toxicity of the monoester metabolite of bis-(2-ethylhexyl) tetrabromophthalate (tbph), a novel brominated flame retardant present in indoor dust. *Environmental health perspectives*, 120(12), 1711–9. doi:10.1289/ehp.1204932
225. Stapleton, H., Klosterhaus, S., Blum, A., & Webster, T. (2010). IDENTIFICATION OF FLAME RETARDANTS IN POLYURETHANE FOAM COLLECTED FROM BABY PRODUCTS. *Organohalogen Compounds*, 72. Retrieved from <http://www.xcdtech.com/dioxin2010/pdf/1310.pdf>
226. Stapleton, H. M., Allen, J. G., Kelly, S. M., Konstantinov, A., Klosterhaus, S., Watkins, D., McClean, M. D., et al. (2008). Alternate and new brominated flame retardants detected in U.S. house dust. *Environmental science & technology*, 42(18), 6910–6. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18853808>

227. Stapleton, H. M., & Dodder, N. G. (2008). Photodegradation of decabromodiphenyl ether in house dust by natural sunlight. *Environmental toxicology and chemistry / SETAC*, 27(2), 306–12. doi:10.1897/07-301R.1
228. Stapleton, H. M., Eagle, S., Anthopolos, R., Wolkin, A., & Miranda, M. L. (2011). Associations between polybrominated diphenyl ether (PBDE) flame retardants, phenolic metabolites, and thyroid hormones during pregnancy. *Environmental health perspectives*, 119(10), 1454–9. doi:10.1289/ehp.1003235
229. Stapleton, H. M., Eagle, S., Sjödin, A., & Webster, T. F. (2012). Serum PBDEs in a North Carolina toddler cohort: associations with handwipes, house dust, and socioeconomic variables. *Environmental health perspectives*, 120(7), 1049–54. doi:10.1289/ehp.1104802
230. Stapleton, H. M., Klosterhaus, S., Eagle, S., Fuh, J., Meeker, J. D., Blum, A., & Webster, T. F. (2009). Detection of organophosphate flame retardants in furniture foam and U.S. house dust. *Environmental science & technology*, 43(19), 7490–5. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2782704&tool=pmcentrez&rendertype=abstract>
231. Stapleton, H. M., Klosterhaus, S., Keller, A., Ferguson, P. L., Van Bergen, S., Cooper, E., Webster, T. F., et al. (2011). Identification of flame retardants in polyurethane foam collected from baby products. *Environmental science & technology*, 45(12), 5323–31. doi:10.1021/es2007462
232. Stapleton, H. M., Sharma, S., Getzinger, G., Ferguson, P. L., Gabriel, M., Webster, T. F., & Blum, A. (2012a). Novel and high volume use flame retardants in US couches reflective of the 2005 PentaBDE phase out. *Environmental science & technology*, 46(24), 13432–9. doi:10.1021/es303471d
233. Stapleton, H. M., Sharma, S., Getzinger, G., Ferguson, P. L., Gabriel, M., Webster, T. F., & Blum, A. (2012b). Novel and high volume use flame retardants in US couches reflective of the 2005 PentaBDE phase out. *Environmental science & technology*, 46(24), 13432–9. doi:10.1021/es303471d
234. Stapleton, H. M., Sjödin, A., Jones, R. S., Niehüser, S., Zhang, Y., & Patterson, D. G. (2008). Serum levels of polybrominated diphenyl ethers (PBDEs) in foam recyclers and carpet installers working in the United States. *Environmental science & technology*, 42(9), 3453–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18522133>
235. Stasinska, A., Reid, A., Hinwood, A., Stevenson, G., Callan, A., Odland, J. Ø., & Heyworth, J. (2013). Concentrations of polybrominated diphenyl ethers (PBDEs) in residential dust samples from Western Australia. *Chemosphere*, 91(2), 187–93. doi:10.1016/j.chemosphere.2012.12.044
236. Stoker, T. E., Cooper, R. L., Lambright, C. S., Wilson, V. S., Furr, J., & Gray, L. E. (2005). In vivo and in vitro anti-androgenic effects of DE-71, a commercial polybrominated diphenyl ether (PBDE) mixture. *Toxicology and applied pharmacology*, 207(1), 78–88. doi:10.1016/j.taap.2005.05.010
237. Sundkvist, A. M., Olofsson, U., & Haglund, P. (2010). Organophosphorus flame retardants and plasticizers in marine and fresh water biota and in human milk. *Journal of environmental monitoring*: JEM, 12(4), 943–51. doi:10.1039/b921910b
238. Suvorov, A., & Takser, L. (2011). Delayed response in the rat frontal lobe transcriptome to perinatal exposure to the flame retardant BDE-47. *Journal of applied toxicology*: JAT, 31(5), 477–83. doi:10.1002/jat.1667
239. Suzuki, G., Nose, K., Takigami, H., Takahashi, S., & Sakai, S. (2006). PBDEs and PBDD/Fs in house and office dust from Japan. *Organohalogen Compounds*, 68, 1843–1846. Retrieved from <http://www.dioxin20xx.org/pdfs/2006/06-425.pdf>
240. Suzuki, Go, Tue, N. M., Malarvannan, G., Sudaryanto, A., Takahashi, S., Tanabe, S., Sakai, S., et al. (2013). Similarities in the endocrine-disrupting potencies of indoor dust and flame retardants by using human osteosarcoma (U2OS) cell-based reporter gene assays. *Environmental science & technology*, 47(6), 2898–908. doi:10.1021/es304691a
241. Szabo, D. T., Richardson, V. M., Ross, D. G., Diliberto, J. J., Kodavanti, P. R. S., & Birnbaum, L. S. (2009). Effects of perinatal PBDE exposure on hepatic phase I, phase II, phase III, and deiodinase 1 gene expression involved in thyroid hormone metabolism in male rat pups. *Toxicological sciences*, 107(1), 27–39. doi:10.1093/toxsci/kfn230

242. Ta, T. A., Koenig, C. M., Golub, M. S., Pessah, I. N., Qi, L., Aronov, P. A., & Berman, R. F. (n.d.). Bioaccumulation and behavioral effects of 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) in perinatally exposed mice. *Neurotoxicology and teratology*, 33(3), 393–404. doi:10.1016/j.ntt.2011.02.003
243. Tagliaferri, S., Caglieri, A., Goldoni, M., Pinelli, S., Alinovi, R., Poli, D., Pellacani, C., et al. (2010). Low concentrations of the brominated flame retardants BDE-47 and BDE-99 induce synergistic oxidative stress-mediated neurotoxicity in human neuroblastoma cells. *Toxicology in vitro: an international journal published in association with BIBRA*, 24(1), 116–22. doi:10.1016/j.tiv.2009.08.020
244. Talsness, C. E., Kuriyama, S. N., Sterner-Kock, A., Schnitker, P., Grande, S. W., Shakibaei, M., Andrade, A., et al. (2008). In utero and lactational exposures to low doses of polybrominated diphenyl ether-47 alter the reproductive system and thyroid gland of female rat offspring. *Environmental health perspectives*, 116(3), 308–14. doi:10.1289/ehp.10536
245. Thomsen, C., Stigum, H., Frøshaug, M., Broadwell, S. L., Becher, G., & Eggesbø, M. (2010). Determinants of brominated flame retardants in breast milk from a large scale Norwegian study. *Environment international*, 36(1), 68–74. doi:10.1016/j.envint.2009.10.002
246. Toms, L.-M. L., Guerra, P., Eljarrat, E., Barceló, D., Harden, F. A., Hobson, P., Sjodin, A., et al. (2012). Brominated flame retardants in the Australian population: 1993-2009. *Chemosphere*, 89(4), 398–403. doi:10.1016/j.chemosphere.2012.05.053
247. Trudel, D., Scheringer, M., Von Goetz, N., & Hungerbühler, K. (2011). Total consumer exposure to polybrominated diphenyl ethers in North America and Europe. *Environmental science & technology*, 45(6), 2391–7. doi:10.1021/es1035046
248. Trudel, D., Tlustos, C., Von Goetz, N., Scheringer, M., & Hungerbühler, K. (n.d.). PBDE exposure from food in Ireland: optimising data exploitation in probabilistic exposure modelling. *Journal of exposure science & environmental epidemiology*, 21(6), 565–75. doi:10.1038/jes.2010.41
249. Trudel, D., Tlustos, C., Von Goetz, N., Scheringer, M., Reichert, P., & Hungerbuhler, K. (2011). Exposure of the Irish population to PBDEs in food: consideration of parameter uncertainty and variability for risk assessment. *Food additives & contaminants. Part A, Chemistry, analysis, control, exposure & risk assessment*, 1–13. doi:10.1080/19440049.2011.572082
250. Tseng, L.-H., Hsu, P.-C., Lee, C.-W., Tsai, S.-S., Pan, M.-H., & Li, M.-H. (2011). Developmental exposure to decabrominated diphenyl ether (BDE-209): Effects on sperm oxidative stress and chromatin dna damage in mouse offspring. *Environmental toxicology*. doi:10.1002/tox.20729
251. Tue, N. M., Takahashi, S., Suzuki, G., Isobe, T., Viet, P. H., Kobara, Y., Seike, N., et al. (2013). Contamination of indoor dust and air by polychlorinated biphenyls and brominated flame retardants and relevance of non-dietary exposure in Vietnamese informal e-waste recycling sites. *Environment international*, 51, 160–7. doi:10.1016/j.envint.2012.11.006
252. Turgeon O'Brien, H., Blanchet, R., Gagné, D., Lauzière, J., Vézina, C., Vaissière, E., Ayotte, P., et al. (2012). Exposure to toxic metals and persistent organic pollutants in Inuit children attending childcare centers in Nunavik, Canada. *Environmental science & technology*, 46(8), 4614–23. doi:10.1021/es203622v
253. Turyk, M. E., Persky, V. W., Imm, P., Knobeloch, L., Chatterton, R., & Anderson, H. A. (2008). Hormone disruption by PBDEs in adult male sport fish consumers. *Environmental health perspectives*, 116(12), 1635–41. doi:10.1289/ehp.11707
254. Vagula, M. C., Kubeldis, N., & Nelatury, C. F. (2011). Effects of BDE-85 on the oxidative status and nerve conduction in rodents. *International journal of toxicology*, 30(4), 428–34. doi:10.1177/1091581811411109
255. Van den Eede, N., Dirtu, A. C., Neels, H., & Covaci, A. (2011). Analytical developments and preliminary assessment of human exposure to organophosphate flame retardants from indoor dust. *Environment international*, 37(2), 454–61. doi:10.1016/j.envint.2010.11.010
256. Van der Veen, I., & De Boer, J. (2012). Phosphorus flame retardants: properties, production, environmental occurrence, toxicity and analysis. *Chemosphere*, 88(10), 1119–53. doi:10.1016/j.chemosphere.2012.03.067

257. Verner, M.-A., Bouchard, M., Fritsche, E., Charbonneau, M., & Haddad, S. (2011). In vitro neurotoxicity data in human risk assessment of polybrominated diphenyl ethers (PBDEs): overview and perspectives. *Toxicology in vitro: an international journal published in association with BIBRA*, 25(8), 1509–15. doi:10.1016/j.tiv.2011.06.007
258. Viberg, H., Fredriksson, A., & Eriksson, P. (2004). Neonatal exposure to the brominated flame-retardant, 2,2',4,4',5-pentabromodiphenyl ether, decreases cholinergic nicotinic receptors in hippocampus and affects spontaneous behaviour in the adult mouse. *Environmental Toxicology and Pharmacology*, 17(2), 61–65. doi:10.1016/j.etap.2004.02.004
259. Viberg, Henrik, & Eriksson, P. (2011). Differences in neonatal neurotoxicity of brominated flame retardants, PBDE 99 and TBBPA, in mice. *Toxicology*, 289(1), 59–65. doi:10.1016/j.tox.2011.07.010
260. Viberg, Henrik, Fredriksson, A., & Eriksson, P. (2003). Neonatal exposure to polybrominated diphenyl ether (PBDE 153) disrupts spontaneous behaviour, impairs learning and memory, and decreases hippocampal cholinergic receptors in adult mice. *Toxicology and applied pharmacology*, 192(2), 95–106. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14550744>
261. Viberg, Henrik, Fredriksson, A., & Eriksson, P. (2007). Changes in spontaneous behaviour and altered response to nicotine in the adult rat, after neonatal exposure to the brominated flame retardant, decabrominated diphenyl ether (PBDE 209). *Neurotoxicology*, 28(1), 136–42. doi:10.1016/j.neuro.2006.08.006
262. Viberg, Henrik, Fredriksson, A., Jakobsson, E., Orn, U., & Eriksson, P. (2003). Neurobehavioral derangements in adult mice receiving decabrominated diphenyl ether (PBDE 209) during a defined period of neonatal brain development. *Toxicological sciences*, 76(1), 112–20. doi:10.1093/toxsci/kfg210
263. Viberg, Henrik, Johansson, N., Fredriksson, A., Eriksson, J., Marsh, G., & Eriksson, P. (2006). Neonatal exposure to higher brominated diphenyl ethers, hepta-, octa-, or nonabromodiphenyl ether, impairs spontaneous behavior and learning and memory functions of adult mice. *Toxicological sciences*, 92(1), 211–8. doi:10.1093/toxsci/kfj196
264. Vizcaino, E., Grimalt, J. O., Lopez-Espinosa, M.-J., Llop, S., Rebagliato, M., & Ballester, F. (2011). Polybromodiphenyl ethers in mothers and their newborns from a non-occupationally exposed population (Valencia, Spain). *Environment international*, 37(1), 152–7. doi:10.1016/j.envint.2010.08.011
265. Vorkamp, K., Thomsen, M., Frederiksen, M., Pedersen, M., & Knudsen, L. E. (2011). Polybrominated diphenyl ethers (PBDEs) in the indoor environment and associations with prenatal exposure. *Environment international*, 37(1), 1–10. doi:10.1016/j.envint.2010.06.001
266. Waaijers, S. L., Kong, D., Hendriks, H. S., De Wit, C. A., Cousins, I. T., Westerink, R. H. S., Leonards, P. E. G., et al. (2013). Persistence, bioaccumulation, and toxicity of halogen-free flame retardants. *Reviews of environmental contamination and toxicology*, 222, 1–71. doi:10.1007/978-1-4614-4717-7_1
267. Wagner, J., Ghosal, S., Whitehead, T., & Metayer, C. (2013). Morphology, spatial distribution, and concentration of flame retardants in consumer products and environmental dusts using scanning electron microscopy and Raman micro-spectroscopy. *Environment international*, 59C, 16–26. doi:10.1016/j.envint.2013.05.003
268. Wang, K.-L., Hsia, S.-M., Mao, I.-F., Chen, M.-L., Wang, S.-W., & Wang, P. S. (2011). Effects of polybrominated diphenyl ethers on steroidogenesis in rat Leydig cells. *Human reproduction (Oxford, England)*, 26(8), 2209–17. doi:10.1093/humrep/der165
269. Wang, L., Zou, W., Zhong, Y., An, J., Zhang, X., Wu, M., & Yu, Z. (2012). The hormesis effect of BDE-47 in HepG2 cells and the potential molecular mechanism. *Toxicology letters*, 209(2), 193–201. doi:10.1016/j.toxlet.2011.12.014
270. Watanabe, W., Shimizu, T., Sawamura, R., Hino, A., Konno, K., & Kurokawa, M. (2010). Functional disorder of primary immunity responding to respiratory syncytial virus infection in offspring mice exposed to a flame retardant, decabrominated diphenyl ether, perinatally. *Journal of medical virology*, 82(6), 1075–82. doi:10.1002/jmv.21770

271. Watkins, D. J., McClean, M. D., Fraser, A. J., Weinberg, J., Stapleton, H. M., Sjödin, A., & Webster, T. F. (2011). Exposure to PBDEs in the office environment: evaluating the relationships between dust, handwipes, and serum. *Environmental health perspectives*, *119*(9), 1247–52. doi:10.1289/ehp.1003271
272. Webster, T., Watkins, D., Walker, C., Fraser, A., Heiger-Bernays, W., Stapleton, H., & McClean, M. (2010). PentaBDE alternatives in homes, offices and cars. *5th International Symposium on Brominated Flame Retardants*. Kyoto, Japan. Retrieved from <http://www.bfr2010.com/abstract-download/2010/90044.pdf>
273. WESCHLER, C., & NAZAROFF, W. (2008). Semivolatile organic compounds in indoor environments. *Atmospheric Environment*, *42*(40), 9018–9040. doi:10.1016/j.atmosenv.2008.09.052
274. Williams, A. L., & DeSesso, J. M. (2010). The potential of selected brominated flame retardants to affect neurological development. *Journal of toxicology and environmental health. Part B, Critical reviews*, *13*(5), 411–48. doi:10.1080/10937401003751630
275. Windham, G. C., Pinney, S. M., Sjödin, A., Lum, R., Jones, R. S., Needham, L. L., Biro, F. M., et al. (2010). Body burdens of brominated flame retardants and other persistent organo-halogenated compounds and their descriptors in US girls. *Environmental research*, *110*(3), 251–7. doi:10.1016/j.envres.2010.01.004
276. Woods, R., Vallero, R. O., Golub, M. S., Suarez, J. K., Ta, T. A., Yasui, D. H., Chi, L.-H., et al. (2012). Long-lived epigenetic interactions between perinatal PBDE exposure and Mecp2308 mutation. *Human molecular genetics*, *21*(11), 2399–411. doi:10.1093/hmg/dd5046
277. Wu, K., Xu, X., Liu, J., Guo, Y., Li, Y., & Huo, X. (2010). Polybrominated diphenyl ethers in umbilical cord blood and relevant factors in neonates from Guiyu, China. *Environmental science & technology*, *44*(2), 813–9. doi:10.1021/es9024518
278. Wu, N., Herrmann, T., Paepke, O., Tickner, J., Hale, R., Harvey, L. E., La Guardia, M., et al. (2007). Human exposure to PBDEs: associations of PBDE body burdens with food consumption and house dust concentrations. *Environmental science & technology*, *41*(5), 1584–9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17396645>
279. Yan, T., Xiang, L., Xuejun, J., Chengzhi, C., Youbin, Q., Xuelan, Y., Yang, L., et al. (2012). Spatial learning and memory deficit of low level polybrominated diphenyl ethers-47 in male adult rat is modulated by intracellular glutamate receptors. *The Journal of toxicological sciences*, *37*(2), 223–33. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22467012>
280. Yu, H.-Y., Zhang, B.-Z., Giesy, J. P., & Zeng, E. Y. (2011). Persistent halogenated compounds in aquaculture environments of South China: implications for global consumers' health risk via fish consumption. *Environment international*, *37*(7), 1190–5. doi:10.1016/j.envint.2011.04.012
281. Zhang, C., Liu, X., & Chen, D. (2010). Role of brominated diphenyl ether-209 in the differentiation of neural stem cells in vitro. *International journal of developmental neuroscience* □: the official journal of the International Society for Developmental Neuroscience, *28*(6), 497–502. doi:10.1016/j.ijdevneu.2010.06.002
282. Zhang, F., Hu, W., Yu, H., Sun, H., Shen, O., Wang, X., Liu, H., et al. (2011). Endocrine disruption effects of 2,2',4,4',6-pentabromodiphenylether (BDE100) in reporter gene assays. *Journal of environmental monitoring* □: JEM, *13*(4), 850–4. doi:10.1039/c0em00654h
283. Zhang, J. G., Sun, X. W., & Ai, H. (2012). Levels and congener profiles of polybrominated diphenyl ethers (PBDEs) in primipara breast milk from Shenzhen and exposure risk for breast-fed infants. *Journal of environmental monitoring* □: JEM, *14*(3), 893–900. doi:10.1039/c2em10739b
284. Zhang, S., Bursian, S. J., Martin, P., Chan, H. M., & Martin, J. (2008). Dietary accumulation, disposition, and metabolism of technical pentabrominated diphenyl ether (DE-71) in pregnant mink (*Mustela vison*) and their offspring. *Environmental toxicology and chemistry / SETAC*, *1*. doi:10.1897/07-487
285. Zhang, W., Cai, Y., Sheng, G., Chen, D., & Fu, J. (2011). Tissue distribution of decabrominated diphenyl ether (BDE-209) and its metabolites in sucking rat pups after prenatal and/or postnatal exposure. *Toxicology*, *283*(1), 49–54. doi:10.1016/j.tox.2011.02.003

286. Zhang, Y., Guo, G. L., Han, X., Zhu, C., Kilfoy, B. A., Zhu, Y., Boyle, P., et al. (2008). Do Polybrominated Diphenyl Ethers (PBDEs) Increase the Risk of Thyroid Cancer? *Bioscience hypotheses*, *1*(4), 195–199. doi:10.1016/j.bihy.2008.06.003
287. Zhao, Y., Ao, H., Chen, L., Sottas, C. M., Ge, R.-S., & Zhang, Y. (2011). Effect of brominated flame retardant BDE-47 on androgen production of adult rat Leydig cells. *Toxicology letters*, *205*(2), 209–14. doi:10.1016/j.toxlet.2011.06.011
288. Zhu, L., Ma, B., & Hites, R. A. (2009). Brominated flame retardants in serum from the general population in northern China. *Environmental science & technology*, *43*(18), 6963–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19806728>
289. Zota, A. R., Adamkiewicz, G., & Morello-Frosch, R. A. (2010). Are PBDEs an environmental equity concern? Exposure disparities by socioeconomic status. *Environmental science & technology*, *44*(15), 5691–2. doi:10.1021/es101723d
290. Zota, A. R., Park, J.-S., Wang, Y., Petreas, M., Zoeller, R. T., & Woodruff, T. J. (2011). Polybrominated diphenyl ethers, hydroxylated polybrominated diphenyl ethers, and measures of thyroid function in second trimester pregnant women in California. *Environmental science & technology*, *45*(18), 7896–905. doi:10.1021/es200422b
291. Zota, A. R., Rudel, R. A., Morello-Frosch, R. A., & Brody, J. G. (2008). Elevated house dust and serum concentrations of PBDEs in California: unintended consequences of furniture flammability standards? *Environmental science & technology*, *42*(21), 8158–64. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19031918>