

Selected Bibliography: Furniture Flame Retardants and Environment

Prepared by Green Science Policy Institute
June 2013

1. Alaee, M. (2003). An overview of commercially used brominated flame retardants, their applications, their use patterns in different countries/regions and possible modes of release. *Environment International*, 29(6), 683–689. doi:10.1016/S0160-4120(03)00121-1
2. Alava, J. J., Lambourn, D., Olesiuk, P., Lance, M., Jeffries, S. J., Gobas, F. A. P. C., & Ross, P. S. (2012). PBDE flame retardants and PCBs in migrating Steller sea lions (*Eumetopias jubatus*) in the Strait of Georgia, British Columbia, Canada. *Chemosphere*, 88(7), 855–64. doi:10.1016/j.chemosphere.2012.03.094
3. Allen, A. (2001). Containment landfills: the myth of sustainability. *Engineering Geology*, 60(1), 3–19. Retrieved from <http://www.ingentaconnect.com/content/els/00137952/2001/00000060/00000001/art00084>
4. Alonso, M. B., Eljarrat, E., Gorga, M., Secchi, E. R., Bassoi, M., Barbosa, L., Bertozzi, C. P., et al. (2012). Natural and anthropogenically-produced brominated compounds in endemic dolphins from Western South Atlantic: another risk to a vulnerable species. *Environmental pollution (Barking, Essex□: 1987)*, 170, 152–60. doi:10.1016/j.envpol.2012.06.001
5. Alvarez, D. A., Maruya, K. A., Dodder, N. G., Lao, W., Furlong, E. T., & Smalling, K. L. (2013). Occurrence of contaminants of emerging concern along the California coast (2009-10) using passive sampling devices. *Marine pollution bulletin*, null(null). doi:10.1016/j.marpolbul.2013.04.022
6. Angioni, S. A., Scorticini, G., Diletti, G., Perletta, F., Ceci, R., & Ferri, N. (n.d.). Study of the toxic effects of flame retardant PBDE-47 on the clam Chamelea gallina (Linnaeus, 1758). *Veterinaria italiana*, 49(1), 69–77. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/23564589>
7. Arkoosh, M. R., Boylen, D., Dietrich, J., Anulacion, B. F., Ginaylitalo, Bravo, C. F., Johnson, L. L., et al. (2010). Disease susceptibility of salmon exposed to polybrominated diphenyl ethers (PBDEs). *Aquatic toxicology*, 98(1), 51–9. doi:10.1016/j.aquatox.2010.01.013
8. Ashizuka, Y., Nakagawa, R., Hori, T., Yasutake, D., Tobiishi, K., & Sasaki, K. (2008). Determination of brominated flame retardants and brominated dioxins in fish collected from three regions of Japan. *Molecular nutrition & food research*, 52(2), 273–83. doi:10.1002/mnfr.200700110
9. Barón, E., Gago-Ferrero, P., Gorga, M., Rudolph, I., Mendoza, G., Zapata, A. M., Díaz-Cruz, S., et al. (2013). Occurrence of hydrophobic organic pollutants (BFRs and UV-filters) in sediments from South America. *Chemosphere*, 92(3), 309–16. doi:10.1016/j.chemosphere.2013.03.032
10. Bartrons, M., Grimalt, J. O., De Mendoza, G., & Catalan, J. (2012). Pollutant dehalogenation capability may depend on the trophic evolutionary history of the organism: PBDEs in freshwater food webs. *PloS one*, 7(7), e41829. doi:10.1371/journal.pone.0041829
11. Beineke, A., Siebert, U., McLachlan, M., Bruhn, R., Thron, K., Failing, K., Müller, G., et al. (2005). Investigations of the potential influence of environmental contaminants on the thymus and spleen of harbor porpoises (*Phocoena phocoena*). *Environmental science & technology*, 39(11), 3933–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15984767>
12. Berg, V., Zerihun, M. A., Jørgensen, A., Lie, E., Dale, O. B., Skaare, J. U., & Lyche, J. L. (2013). High prevalence of infections and pathological changes in burbot (*Lota lota*) from a polluted lake (Lake Mjøsa, Norway). *Chemosphere*, 90(5), 1711–8. doi:10.1016/j.chemosphere.2012.10.017
13. Besis, A., & Samara, C. (2012). Polybrominated diphenyl ethers (PBDEs) in the indoor and outdoor environments--a review on occurrence and human exposure. *Environmental pollution (Barking, Essex□: 1987)*, 169, 217–29. doi:10.1016/j.envpol.2012.04.009
14. Birnbaum, L. S., & Staskal, D. F. (2004). Brominated flame retardants: cause for concern? *Environmental health perspectives*, 112(1), 9–17. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1241790&tool=pmcentrez&rendertype=abstract>

15. Björklund, K., Strömvall, A.-M., & Malmqvist, P.-A. (2011). Screening of organic contaminants in urban snow. *Water science and technology: a journal of the International Association on Water Pollution Research*, 64(1), 206–13. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22053476>
16. Brambilla, G., De Filippis, S. P., Iamiceli, A. L., Iacovella, N., Abate, V., Aronica, V., Di Marco, V., et al. (2011). Bioaccumulation of dioxin-like substances and selected brominated flame retardant congeners in the fat and livers of black pigs farmed within the Nebrodi Regional Park of Sicily. *Journal of food protection*, 74(2), 261–9. doi:10.4315/0362-028X.JFP-10-176
17. Breivik, K., Gioia, R., Chakraborty, P., Zhang, G., & Jones, K. C. (2011). Are reductions in industrial organic contaminants emissions in rich countries achieved partly by export of toxic wastes? *Environmental science & technology*, 45(21), 9154–60. doi:10.1021/es202320c
18. Brown, T. N., & Wania, F. (2008). Screening chemicals for the potential to be persistent organic pollutants: a case study of Arctic contaminants. *Environmental science & technology*, 42(14), 5202–9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18754370>
19. Cao, S., Zeng, X., Song, H., Li, H., Yu, Z., Sheng, G., & Fu, J. (2012). Levels and distributions of organophosphate flame retardants and plasticizers in sediment from Taihu Lake, China. *Environmental toxicology and chemistry / SETAC*, 31(7), 1478–84. doi:10.1002/etc.1872
20. Carlsson, P., Herzke, D., Wedborg, M., & Gabrielsen, G. W. (2011). Environmental pollutants in the Swedish marine ecosystem, with special emphasis on polybrominated diphenyl ethers (PBDE). *Chemosphere*, 82(9), 1286–92. doi:10.1016/j.chemosphere.2010.12.029
21. Carpenter, D. O. (2011). Health effects of persistent organic pollutants: the challenge for the Pacific Basin and for the world. *Reviews on environmental health*, 26(1), 61–9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21714383>
22. Cary, T. L., & Karasov, W. H. (2013). Toxicokinetics of polybrominated diphenyl ethers across life stages in the northern leopard frog (*Lithobates pipiens*). *Environmental toxicology and chemistry / SETAC*, 32(7), 1631–40. doi:10.1002/etc.2215
23. Chabot-Giguère, B., Letcher, R. J., & Verreault, J. (2013). In vitro biotransformation of decabromodiphenyl ether (BDE-209) and Dechlorane Plus flame retardants: a case study of ring-billed gull breeding in a pollution hotspot in the St. Lawrence River, Canada. *Environment international*, 55, 101–8. doi:10.1016/j.envint.2013.02.008
24. Champoux, L., Moisey, J., & Muir, D. C. G. (2010). Polybrominated diphenyl ethers, toxaphenes, and other halogenated organic pollutants in great blue heron eggs. *Environmental toxicology and chemistry / SETAC*, 29(2), 243–9. doi:10.1002/etc.37
25. Chan, W. K., & Chan, K. M. (2012). Disruption of the hypothalamic-pituitary-thyroid axis in zebrafish embryos-larvae following waterborne exposure to BDE-47, TBBPA and BPA. *Aquatic toxicology (Amsterdam, Netherlands)*, 108, 106–11. doi:10.1016/j.aquatox.2011.10.013
26. Chen, D., & Hale, R. C. (2010). A global review of polybrominated diphenyl ether flame retardant contamination in birds. *Environment international*, 36(7), 800–11. doi:10.1016/j.envint.2010.05.013
27. Chen, D., Hale, R. C., Watts, B. D., La Guardia, M. J., Harvey, E., & Mojica, E. K. (2010). Species-specific accumulation of polybrominated diphenyl ether flame retardants in birds of prey from the Chesapeake Bay region, USA. *Environmental pollution (Barking, Essex: 1987)*, 158(5), 1883–9. doi:10.1016/j.envpol.2009.10.042
28. Chen, D., Letcher, R. J., Burgess, N. M., Champoux, L., Elliott, J. E., Hebert, C. E., Martin, P., et al. (2012). Flame retardants in eggs of four gull species (Laridae) from breeding sites spanning Atlantic to Pacific Canada. *Environmental pollution (Barking, Essex: 1987)*, 168, 1–9. doi:10.1016/j.envpol.2012.03.040
29. Chen, D., Letcher, R. J., & Chu, S. (2012). Determination of non-halogenated, chlorinated and brominated organophosphate flame retardants in herring gull eggs based on liquid chromatography-tandem quadrupole mass spectrometry. *Journal of chromatography. A*, 1220, 169–74. doi:10.1016/j.chroma.2011.11.046

30. Chen, D., Letcher, R. J., & Martin, P. (2012). Flame retardants in eggs of American kestrels and European starlings from southern Lake Ontario region (North America). *Journal of environmental monitoring* □: JEM, 14(11), 2870–6. doi:10.1039/c2em30472d
31. Chen, S.-J., Feng, A.-H., He, M.-J., Chen, M.-Y., Luo, X.-J., & Mai, B.-X. (2013). Current levels and composition profiles of PBDEs and alternative flame retardants in surface sediments from the Pearl River Delta, southern China: comparison with historical data. *The Science of the total environment*, 444, 205–11. doi:10.1016/j.scitotenv.2012.11.104
32. Chen, X., Huang, C., Wang, X., Chen, J., Bai, C., Chen, Y., Chen, X., et al. (2012). BDE-47 disrupts axonal growth and motor behavior in developing zebrafish. *Aquatic toxicology (Amsterdam, Netherlands)*, 120-121, 35–44. doi:10.1016/j.aquatox.2012.04.014
33. Chiu, J. M. Y., Po, B. H. K., Chan, C. Y. S., Lam, M. H. W., Qian, P.-Y., & Wu, R. S. S. (2012). Polybrominated diphenyl ethers (PBDEs) alter larval settlement of marine intertidal organisms across three phyla via reducing bacterial abundance on the biofilms. *Environmental science & technology*, 46(14), 7772–81. doi:10.1021/es300261c
34. Chiu, K. H., Lin, C.-R., Huang, H.-W., Shiea, J., & Liu, L. L. (2012). Toxic effects of two brominated flame retardants BDE-47 and BDE-183 on the survival and protein expression of the tubificid *Monopylephorus limosus*. *Ecotoxicology and environmental safety*, 84, 46–53. doi:10.1016/j.ecoenv.2012.06.022
35. Chou, C.-T., Hsiao, Y.-C., Ko, F.-C., Cheng, J.-O., Cheng, Y.-M., & Chen, T.-H. (2010). Chronic exposure of 2,2',4,4'-tetrabromodiphenyl ether (PBDE-47) alters locomotion behavior in juvenile zebrafish (*Danio rerio*). *Aquatic toxicology*, 98(4), 388–95. doi:10.1016/j.aquatox.2010.03.012
36. Cincinelli, A., Martellini, T., Misuri, L., Lanciotti, E., Sweetman, A., Laschi, S., & Palchetti, I. (2012). PBDEs in Italian sewage sludge and environmental risk of using sewage sludge for land application. *Environmental pollution (Barking, Essex□: 1987)*, 161, 229–34. doi:10.1016/j.envpol.2011.11.001
37. Covaci, A., Harrad, S., Abdallah, M. A.-E., Ali, N., Law, R. J., Herzke, D., & De Wit, C. A. (2011). Novel brominated flame retardants: a review of their analysis, environmental fate and behaviour. *Environment international*, 37(2), 532–56. doi:10.1016/j.envint.2010.11.007
38. Cristale, J., Katsoyiannis, A., Sweetman, A. J., Jones, K. C., & Lacorte, S. (2013). Occurrence and risk assessment of organophosphorus and brominated flame retardants in the River Aire (UK). *Environmental pollution (Barking, Essex□: 1987)*, 179C(null), 194–200. doi:10.1016/j.envpol.2013.04.001
39. Crump, D., Chiu, S., & Kennedy, S. W. (2012). Effects of tris(1,3-dichloro-2-propyl) phosphate and tris(1-chloropropyl) phosphate on cytotoxicity and mRNA expression in primary cultures of avian hepatocytes and neuronal cells. *Toxicological sciences□: an official journal of the Society of Toxicology*, 126(1), 140–8. doi:10.1093/toxsci/kfs015
40. Daso, A. P., Fatoki, O. S., Odendaal, J. P., & Olujimi, O. O. (2012). Occurrence of selected polybrominated diphenyl ethers and 2,2',4,4',5,5'-hexabromobiphenyl (BB-153) in sewage sludge and effluent samples of a wastewater-treatment plant in Cape Town, South Africa. *Archives of environmental contamination and toxicology*, 62(3), 391–402. doi:10.1007/s00244-011-9720-9
41. Davies, R., & Zou, E. (2012). Polybrominated diphenyl ethers disrupt molting in neonatal *Daphnia magna*. *Ecotoxicology (London, England)*, 21(5), 1371–80. doi:10.1007/s10646-012-0891-6
42. Davis, E. F., Klosterhaus, S. L., & Stapleton, H. M. (2012). Measurement of flame retardants and triclosan in municipal sewage sludge and biosolids. *Environment international*, 40, 1–7. doi:10.1016/j.envint.2011.11.008
43. De Wit, C. A. (2002a). An overview of brominated flame retardants in the environment. *Chemosphere*, 46(5), 583–624. doi:10.1016/S0045-6535(01)00225-9
44. De Wit, C. A. (2002b). An overview of brominated flame retardants in the environment. *Chemosphere*, 46(5), 583–624. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11999784>
45. De Wit, C. A., Herzke, D., & Vorkamp, K. (2009). Brominated flame retardants in the Arctic environment - trends and new candidates. *The Science of the total environment*, 408(15), 2885–918. doi:10.1016/j.scitotenv.2009.08.037

46. Desforges, J.-P. W., Ross, P. S., & Loseto, L. L. (2012). Transplacental transfer of polychlorinated biphenyls and polybrominated diphenyl ethers in arctic beluga whales (*Delphinapterus leucas*). *Environmental toxicology and chemistry / SETAC*, 31(2), 296–300. doi:10.1002/etc.750
47. Dietz, R., Rigét, F. F., Sonne, C., Born, E. W., Bechshøft, T., McKinney, M. A., Drimmie, R. J., et al. (2012). Three decades (1983–2010) of contaminant trends in East Greenland polar bears (*Ursus maritimus*). Part 2: Brominated flame retardants. *Environment international*. doi:10.1016/j.envint.2012.09.008
48. Dorneles, P. R., Lailson-Brito, J., Dirtu, A. C., Weijs, L., Azevedo, A. F., Torres, J. P. M., Malm, O., et al. (2010). Anthropogenic and naturally-produced organobrominated compounds in marine mammals from Brazil. *Environment international*, 36(1), 60–7. doi:10.1016/j.envint.2009.10.001
49. Duan, Y.-P., Meng, X.-Z., Yang, C., Pan, Z.-Y., Chen, L., Yu, R., & Li, F.-T. (2010). Polybrominated diphenyl ethers in background surface soils from the Yangtze River Delta (YRD), China: occurrence, sources, and inventory. *Environmental science and pollution research international*, 17(4), 948–56. doi:10.1007/s11356-010-0295-1
50. Eggen, T., Heimstad, E. S., Stuanes, A. O., & Norli, H. R. (2012). Uptake and translocation of organophosphates and other emerging contaminants in food and forage crops. *Environmental science and pollution research international*. doi:10.1007/s11356-012-1363-5
51. Eguchi, A., Isobe, T., Ramu, K., Tue, N. M., Sudaryanto, A., Devanathan, G., Viet, P. H., et al. (2013). Soil contamination by brominated flame retardants in open waste dumping sites in Asian developing countries. *Chemosphere*, 90(9), 2365–71. doi:10.1016/j.chemosphere.2012.10.027
52. Eisenreich, K. M., & Rowe, C. L. (2013). Experimental exposure of eggs to polybrominated diphenyl ethers BDE-47 and BDE-99 in red-eared sliders (*Trachemys scripta elegans*) and snapping turtles (*Chelydra serpentina*) and possible species-specific differences in debromination. *Environmental toxicology and chemistry / SETAC*, 32(2), 393–400. doi:10.1002/etc.2061
53. Elliott, J. E., Wilson, L. K., & Wakeford, B. (2005). Polybrominated diphenyl ether trends in eggs of marine and freshwater birds from British Columbia, Canada, 1979–2002. *Environmental science & technology*, 39(15), 5584–91. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16124290>
54. Eng, M. L., Elliott, J. E., MacDougall-Shackleton, S. A., Letcher, R. J., & Williams, T. D. (2012). Early exposure to 2,2',4,4',5-pentabromodiphenyl ether (BDE-99) affects mating behavior of zebra finches. *Toxicological sciences: an official journal of the Society of Toxicology*, 127(1), 269–76. doi:10.1093/toxsci/kfs076
55. Eng, M. L., Williams, T. D., & Elliott, J. E. (2013). Developmental exposure to a brominated flame retardant: An assessment of effects on physiology, growth, and reproduction in a songbird, the zebra finch. *Environmental pollution (Barking, Essex): 1987)*, 178(null), 343–9. doi:10.1016/j.envpol.2013.03.037
56. Fair, P. A., Mitchum, G., Hulsey, T. C., Adams, J., Zolman, E., McFee, W., Wirth, E., et al. (2007). Polybrominated diphenyl ethers (PBDEs) in blubber of free-ranging bottlenose dolphins (*Tursiops truncatus*) from two southeast Atlantic estuarine areas. *Archives of environmental contamination and toxicology*, 53(3), 483–94. doi:10.1007/s00244-006-0244-7
57. Farhat, A., Crump, D., Chiu, S., Williams, K. L., Letcher, R. J., Gauthier, L. T., & Kennedy, S. W. (2013). In Ovo Effects of Two Organophosphate Flame Retardants--TCPP and TDCPP--on Pipping Success, Development, mRNA Expression, and Thyroid Hormone Levels in Chicken Embryos. *Toxicological sciences: an official journal of the Society of Toxicology*. doi:10.1093/toxsci/kft100
58. Feng, C., Xu, Y., Zhao, G., Zha, J., Wu, F., & Wang, Z. (2012). Relationship between BDE 209 metabolites and thyroid hormone levels in rainbow trout (*Oncorhynchus mykiss*). *Aquatic toxicology (Amsterdam, Netherlands)*, 122-123, 28–35. doi:10.1016/j.aquatox.2012.05.008
59. Fernie, K. J., & Letcher, R. J. (2010). Historical contaminants, flame retardants, and halogenated phenolic compounds in peregrine Falcon (*Falco peregrinus*) nestlings in the Canadian Great Lakes Basin. *Environmental science & technology*, 44(9), 3520–6. doi:10.1021/es100400n

60. Fernie, K. J., Shutt, J. L., Letcher, R. J., Ritchie, I. J., & Bird, D. M. (2009). Environmentally relevant concentrations of DE-71 and HBCD alter eggshell thickness and reproductive success of American kestrels. *Environmental science & technology*, 43(6), 2124–30. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19368224>
61. Fernie, K. J., Shutt, J. L., Letcher, R. J., Ritchie, I. J., Sullivan, K., & Bird, D. M. (2008). Changes in reproductive courtship behaviors of adult American kestrels (*Falco sparverius*) exposed to environmentally relevant levels of the polybrominated diphenyl ether mixture, DE-71. *Toxicological sciences*, 102(1), 171–8. doi:10.1093/toxsci/kfm295
62. Fernie, K. J., Shutt, J. L., Mayne, G., Hoffman, D., Letcher, R. J., Drouillard, K. G., & Ritchie, I. J. (2005). Exposure to polybrominated diphenyl ethers (PBDEs): changes in thyroid, vitamin A, glutathione homeostasis, and oxidative stress in American kestrels (*Falco sparverius*). *Toxicological sciences*, 88(2), 375–83. doi:10.1093/toxsci/kfi295
63. Fries, E., & Mihajlović, I. (2011). Pollution of soils with organophosphorus flame retardants and plasticizers. *Journal of environmental monitoring* □: JEM, 13(10), 2692–4. doi:10.1039/c1em10538h
64. Frouin, H., Lebeuf, M., Hammill, M., Sjare, B., & Fournier, M. (2011). PBDEs in serum and blubber of harbor, grey and harp seal pups from Eastern Canada. *Chemosphere*, 82(5), 663–9. doi:10.1016/j.chemosphere.2010.11.007
65. Fujii, Y., Ito, Y., Harada, K. H., Hitomi, T., Koizumi, A., & Haraguchi, K. (2012). Regional variation and possible sources of brominated contaminants in breast milk from Japan. *Environmental pollution (Barking, Essex)* □: 1987, 162, 269–74. doi:10.1016/j.envpol.2011.11.022
66. Gao, F., Luo, X.-J., Yang, Z.-F., Wang, X.-M., & Mai, B.-X. (2009). Brominated flame retardants, polychlorinated biphenyls, and organochlorine pesticides in bird eggs from the Yellow River Delta, North China. *Environmental science & technology*, 43(18), 6956–62. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19806727>
67. Gaylor, M. O., Harvey, E., & Hale, R. C. (2012). House crickets can accumulate polybrominated diphenyl ethers (PBDEs) directly from polyurethane foam common in consumer products. *Chemosphere*, 86(5), 500–5. doi:10.1016/j.chemosphere.2011.10.014
68. Gentes, M.-L., Letcher, R. J., Caron-Beaudoin, E., & Verreault, J. (2012). Novel flame retardants in urban-feeding ring-billed gulls from the St. Lawrence River, Canada. *Environmental science & technology*, 46(17), 9735–44. doi:10.1021/es302099f
69. Gorga, M., Martínez, E., Ginebreda, A., Eljarrat, E., & Barceló, D. (2013). Determination of PBDEs, HBB, PBEB, DBDPE, HBCD, TBBPA and related compounds in sewage sludge from Catalonia (Spain). *The Science of the total environment*, 444, 51–9. doi:10.1016/j.scitotenv.2012.11.066
70. Grenier, J. L., & Davis, J. A. (2010). Water quality in South San Francisco Bay, California: current condition and potential issues for the South Bay Salt Pond Restoration Project. *Reviews of environmental contamination and toxicology*, 206, 115–47. doi:10.1007/978-1-4419-6260-7_6
71. Guerra, P., Alaee, M., Jiménez, B., Pacepavicius, G., Marvin, C., MacInnis, G., Eljarrat, E., et al. (2012). Emerging and historical brominated flame retardants in peregrine falcon (*Falco peregrinus*) eggs from Canada and Spain. *Environment international*, 40, 179–86. doi:10.1016/j.envint.2011.07.014
72. Hale, R. (2003). Polybrominated diphenyl ether flame retardants in the North American environment. *Environment International*, 29(6), 771–779. doi:10.1016/S0160-4120(03)00113-2
73. Hale, R. C., La Guardia, M. J., Harvey, E., Chen, D., Mainor, T. M., Luellen, D. R., & Hundal, L. S. (2012). Polybrominated diphenyl ethers in U.S. sewage sludges and biosolids: temporal and geographical trends and uptake by corn following land application. *Environmental science & technology*, 46(4), 2055–63. doi:10.1021/es203149g
74. Hale, R. C., La Guardia, M. J., Harvey, E., Gaylor, M. O., & Mainor, T. M. (2006). Brominated flame retardant concentrations and trends in abiotic media. *Chemosphere*, 64(2), 181–6. doi:10.1016/j.chemosphere.2005.12.006

75. Hall, A. J., Kalantzi, O. I., & Thomas, G. O. (2003). Polybrominated diphenyl ethers (PBDEs) in grey seals during their first year of life--are they thyroid hormone endocrine disrupters? *Environmental pollution*, 126(1), 29–37. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12860100>
76. Hall, A. J., & Thomas, G. O. (2007). Polychlorinated biphenyls, DDT, polybrominated diphenyl ethers, and organic pesticides in United Kingdom harbor seals (*Phoca vitulina*)--mixed exposures and thyroid homeostasis. *Environmental toxicology and chemistry / SETAC*, 26(5), 851–61. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17521129>
77. Hall, A. J., Thomas, G. O., & McConnell, B. J. (2009). Exposure to persistent organic pollutants and first-year survival probability in gray seal pups. *Environmental science & technology*, 43(16), 6364–9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19746738>
78. He, J., Yang, D., Wang, C., Liu, W., Liao, J., Xu, T., Bai, C., et al. (2011). Chronic zebrafish low dose decabrominated diphenyl ether (BDE-209) exposure affected parental gonad development and locomotion in F1 offspring. *Ecotoxicology (London, England)*, 20(8), 1813–22. doi:10.1007/s10646-011-0720-3
79. He, M.-J., Luo, X.-J., Chen, M.-Y., Sun, Y.-X., Chen, S.-J., & Mai, B.-X. (2012). Bioaccumulation of polybrominated diphenyl ethers and decabromodiphenyl ethane in fish from a river system in a highly industrialized area, South China. *The Science of the total environment*, 419, 109–15. doi:10.1016/j.scitotenv.2011.12.035
80. Henny, C. J., Kaiser, J. L., Grove, R. A., Johnson, B. L., & Letcher, R. J. (2009). Polybrominated diphenyl ether flame retardants in eggs may reduce reproductive success of ospreys in Oregon and Washington, USA. *Ecotoxicology*, 18(7), 802–13. doi:10.1007/s10646-009-0323-4
81. Hermanson, M. H., Isaksson, E., Forsström, S., Teixeira, C., Muir, D. C. G., Pohjola, V. A., & Van de Wal, R. S. V. (2010). Deposition history of brominated flame retardant compounds in an ice core from Holtedahlfonna, Svalbard, Norway. *Environmental science & technology*, 44(19), 7405–10. doi:10.1021/es1016608
82. Hloušková, V., Lanková, D., Kalachová, K., Hrádková, P., Pousta, J., Hajšlová, J., & Pulkrabová, J. (2013). Occurrence of brominated flame retardants and perfluoroalkyl substances in fish from the Czech aquatic ecosystem. *The Science of the total environment*, 461-462C, 88–98. doi:10.1016/j.scitotenv.2013.04.081
83. Hrádková, P., Pulkrabová, J., Kalachová, K., Hloušková, V., Tomaniová, M., Pousta, J., & Hajšlová, J. (2012). Occurrence of halogenated contaminants in fish from selected river localities and ponds in the Czech Republic. *Archives of environmental contamination and toxicology*, 62(1), 85–96. doi:10.1007/s00244-011-9681-z
84. Hu, G.-C., Luo, X.-J., Dai, J.-Y., Zhang, X.-L., Wu, H., Zhang, C.-L., Guo, W., et al. (2008). Brominated flame retardants, polychlorinated biphenyls, and organochlorine pesticides in captive giant panda (*Ailuropoda melanoleuca*) and red panda (*Ailurus fulgens*) from China. *Environmental science & technology*, 42(13), 4704–9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18677994>
85. Huang, H., Zhang, S., Christie, P., Wang, S., & Xie, M. (2010). Behavior of decabromodiphenyl ether (BDE-209) in the soil-plant system: uptake, translocation, and metabolism in plants and dissipation in soil. *Environmental science & technology*, 44(2), 663–7. doi:10.1021/es901860r
86. Ikonomou, M. G., Rayne, S., & Addison, R. F. (2002). Exponential increases of the brominated flame retardants, polybrominated diphenyl ethers, in the Canadian Arctic from 1981 to 2000. *Environmental science & technology*, 36(9), 1886–92. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12026966>
87. Ilyas, M., Sudaryanto, A., Setiawan, I. E., Riyadi, A. S., Isobe, T., Ogawa, S., Takahashi, S., et al. (2011). Characterization of polychlorinated biphenyls and brominated flame retardants in surface soils from Surabaya, Indonesia. *Chemosphere*, 83(6), 783–91. doi:10.1016/j.chemosphere.2011.02.067
88. Ilyas, M., Sudaryanto, A., Setiawan, I. E., Riyadi, A. S., Isobe, T., Takahashi, S., & Tanabe, S. (2011). Characterization of polychlorinated biphenyls and brominated flame retardants in sediments from riverine and coastal waters of Surabaya, Indonesia. *Marine pollution bulletin*, 62(1), 89–98. doi:10.1016/j.marpolbul.2010.09.006
89. Isobe, T., Oshihori, T., Hamada, H., Nakayama, K., Yamada, T. K., Tajima, Y., Amano, M., et al. (2011). Contamination status of POPs and BFRs and relationship with parasitic infection in finless porpoises

- (*Neophocaena phocaenoides*) from Seto Inland Sea and Omura Bay, Japan. *Marine pollution bulletin*, 63(5-12), 564–71. doi:10.1016/j.marpolbul.2011.01.014
90. Johansson, A.-K., Sellström, U., Lindberg, P., Bignert, A., & De Wit, C. A. (2011). Temporal trends of polybrominated diphenyl ethers and hexabromocyclododecane in Swedish Peregrine Falcon (*Falco peregrinus peregrinus*) eggs. *Environment international*, 37(4), 678–86. doi:10.1016/j.envint.2011.01.010
91. Johansson, A.-K., Sellström, U., Lindberg, P., Bignert, A., & De Witt, C. A. (2009). Polybrominated diphenyl ether congener patterns, hexabromocyclododecane, and brominated biphenyl 153 in eggs of peregrine falcons (*Falco peregrinus*) breeding in Sweden. *Environmental toxicology and chemistry / SETAC*, 28(1), 9–17. doi:10.1897/08-142.1
92. Johnson-Restrepo, B., Kannan, K., Addink, R., & Adams, D. H. (2005). Polybrominated diphenyl ethers and polychlorinated biphenyls in a marine foodweb of coastal Florida. *Environmental science & technology*, 39(21), 8243–50. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16294860>
93. Kannan, K., Perrotta, E., Thomas, N. J., & Aldous, K. M. (2007). A comparative analysis of polybrominated diphenyl ethers and polychlorinated biphenyls in Southern sea otters that died of infectious diseases and noninfectious causes. *Archives of environmental contamination and toxicology*, 53(2), 293–302. doi:10.1007/s00244-006-0251-8
94. Karlsson, M., Ericson, I., Van Bavel, B., Jensen, J.-K., & Dam, M. (2006). Levels of brominated flame retardants in Northern Fulmar (*Fulmarus glacialis*) eggs from the Faroe Islands. *The Science of the total environment*, 367(2-3), 840–6. doi:10.1016/j.scitotenv.2006.02.050
95. Klecka, G., Persoon, C., & Currie, R. (2010). Chemicals of emerging concern in the Great Lakes Basin: an analysis of environmental exposures. *Reviews of environmental contamination and toxicology*, 207, 1–93. doi:10.1007/978-1-4419-6406-9_1
96. Klosterhaus, S. L., Dreis, E., & Baker, J. E. (2011). Bioaccumulation kinetics of polybrominated diphenyl ethers from estuarine sediments to the marine polychaete, *Nereis virens*. *Environmental toxicology and chemistry / SETAC*, 30(5), 1204–12. doi:10.1002/etc.497
97. Klosterhaus, S. L., Stapleton, H. M., La Guardia, M. J., & Greig, D. J. (2012). Brominated and chlorinated flame retardants in San Francisco Bay sediments and wildlife. *Environment international*, 47, 56–65. doi:10.1016/j.envint.2012.06.005
98. Krahn, M. M., Hanson, M. B., Baird, R. W., Boyer, R. H., Burrows, D. G., Emmons, C. K., Ford, J. K. B., et al. (2007). Persistent organic pollutants and stable isotopes in biopsy samples (2004/2006) from Southern Resident killer whales. *Marine pollution bulletin*, 54(12), 1903–11. doi:10.1016/j.marpolbul.2007.08.015
99. Kucklick, J., Schwacke, L., Wells, R., Hohn, A., Guichard, A., Yordy, J., Hansen, L., et al. (2011). Bottlenose dolphins as indicators of persistent organic pollutants in the western North Atlantic Ocean and northern Gulf of Mexico. *Environmental science & technology*, 45(10), 4270–7. doi:10.1021/es1042244
100. Kuo, Y.-M., Sepúlveda, M. S., Hua, I., Ochoa-Acuña, H. G., & Sutton, T. M. (2010). Bioaccumulation and biomagnification of polybrominated diphenyl ethers in a food web of Lake Michigan. *Ecotoxicology (London, England)*, 19(4), 623–34. doi:10.1007/s10646-009-0431-1
101. Kuo, Y.-M., Sepúlveda, M. S., Sutton, T. M., Ochoa-Acuña, H. G., Muir, A. M., Miller, B., & Hua, I. (2010). Bioaccumulation and biotransformation of decabromodiphenyl ether and effects on daily growth in juvenile lake whitefish (*Coregonus clupeaformis*). *Ecotoxicology (London, England)*, 19(4), 751–60. doi:10.1007/s10646-009-0451-x
102. Kwan, C. S., Takada, H., Mizukawa, K., Torii, M., Koike, T., Yamashita, R., Rinawati, et al. (2013). PBDEs in leachates from municipal solid waste dumping sites in tropical Asian countries: phase distribution and debromination. *Environmental science and pollution research international*, 20(6), 4188–204. doi:10.1007/s11356-012-1365-3
103. La Guardia, M. J., Hale, R. C., & Harvey, E. (2007). Evidence of debromination of decabromodiphenyl ether (BDE-209) in biota from a wastewater receiving stream. *Environmental science & technology*, 41(19), 6663–70. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17969678>

- 104.La Guardia, M. J., Hale, R. C., Harvey, E., & Chen, D. (2010). Flame-retardants and other organohalogens detected in sewage sludge by electron capture negative ion mass spectrometry. *Environmental science & technology*, 44(12), 4658–64. doi:10.1021/es9039264
- 105.La Guardia, M. J., Hale, R. C., Harvey, E., Mainor, T. M., & Ciparis, S. (2012). In situ accumulation of HBCD, PBDEs, and several alternative flame-retardants in the bivalve (*Corbicula fluminea*) and gastropod (*Elimia proxima*). *Environmental science & technology*, 46(11), 5798–805. doi:10.1021/es3004238
- 106.Lam, J. C. W., Lau, R. K. F., Murphy, M. B., & Lam, P. K. S. (2009). Temporal trends of hexabromocyclododecanes (HBCDs) and polybrominated diphenyl ethers (PBDEs) and detection of two novel flame retardants in marine mammals from Hong Kong, South China. *Environmental science & technology*, 43(18), 6944–9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19806725>
- 107.Laniewski, K., BorÉn, H., & Grimvall, A. (1998). Identification of Volatile and Extractable Chloroorganics in Rain and Snow. *Environmental Science & Technology*, 32(24), 3935–3940. doi:10.1021/es980480i
- 108.Law, K., Halldorson, T., Danell, R., Stern, G., Gewurtz, S., Alaee, M., Marvin, C., et al. (2006). Bioaccumulation and trophic transfer of some brominated flame retardants in a Lake Winnipeg (Canada) food web. *Environmental toxicology and chemistry / SETAC*, 25(8), 2177–86. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16916037>
- 109.Law, R. J., Allchin, C. R., De Boer, J., Covaci, A., Herzke, D., Lepom, P., Morris, S., et al. (2006). Levels and trends of brominated flame retardants in the European environment. *Chemosphere*, 64(2), 187–208. doi:10.1016/j.chemosphere.2005.12.007
- 110.Lee, H. J., Kim, G. B., & Lee, R. F. (2012). Genotoxicity and development effects of brominated flame retardant PBDEs and UV-exposed PBDEs on grass shrimp (*Palaemonetes pugio*) embryo. *Marine pollution bulletin*, 64(12), 2892–5. doi:10.1016/j.marpolbul.2012.08.010
- 111.Lehner, A. F., Samsing, F., & Rumbeiha, W. K. (2010). Organophosphate ester flame retardant-induced acute intoxications in dogs. *Journal of medical toxicology*: official journal of the American College of Medical Toxicology, 6(4), 448–58. doi:10.1007/s13181-010-0105-7
- 112.Lema, S. C., Schultz, I. R., Scholz, N. L., Incardona, J. P., & Swanson, P. (2007). Neural defects and cardiac arrhythmia in fish larvae following embryonic exposure to 2,2',4,4'-tetrabromodiphenyl ether 47. *Aquatic toxicology*, 82(4), 296–307. doi:10.1016/j.aquatox.2007.03.002
- 113.Letcher, R. J., Bustnes, J. O., Dietz, R., Jenssen, B. M., Jørgensen, E. H., Sonne, C., Verreault, J., et al. (2010). Exposure and effects assessment of persistent organohalogen contaminants in arctic wildlife and fish. *The Science of the total environment*, 408(15), 2995–3043. doi:10.1016/j.scitotenv.2009.10.038
- 114.Letcher, R. J., Gebbink, W. A., Sonne, C., Born, E. W., McKinney, M. A., & Dietz, R. (2009). Bioaccumulation and biotransformation of brominated and chlorinated contaminants and their metabolites in ringed seals (*Pusa hispida*) and polar bears (*Ursus maritimus*) from East Greenland. *Environment international*, 35(8), 1118–24. doi:10.1016/j.envint.2009.07.006
- 115.Li, Juan, Liang, Y., Zhang, X., Lu, J., Zhang, J., Ruan, T., Zhou, Q., et al. (2011). Impaired gas bladder inflation in zebrafish exposed to a novel heterocyclic brominated flame retardant tris(2,3-dibromopropyl) isocyanurate. *Environmental science & technology*, 45(22), 9750–7. doi:10.1021/es202420g
- 116.Li, Jun, Zhang, G., Xu, Y., Liu, X., & Li, X.-D. (2010). Dry and wet particle deposition of polybrominated diphenyl ethers (PBDEs) in Guangzhou and Hong Kong, South China. *Journal of environmental monitoring*: JEM, 12(9), 1730–6. doi:10.1039/c001526a
- 117.Li, W., Zhu, L., Zha, J., & Wang, Z. (2011). Effects of decabromodiphenyl ether (BDE-209) on mRNA transcription of thyroid hormone pathway and spermatogenesis associated genes in Chinese rare minnow (*Gobiocypris rarus*). *Environmental toxicology*. doi:10.1002/tox.20767
- 118.Liu, C., Wang, Q., Liang, K., Liu, J., Zhou, B., Zhang, X., Liu, H., et al. (2013). Effects of tris(1,3-dichloro-2-propyl) phosphate and triphenyl phosphate on receptor-associated mRNA expression in zebrafish embryos/larvae. *Aquatic toxicology (Amsterdam, Netherlands)*, 128-129, 147–57. doi:10.1016/j.aquatox.2012.12.010

- 119.Liu, P.-Y., Du, G.-D., Zhao, Y.-X., Mu, Y.-S., Zhang, A.-Q., Qin, Z.-F., Zhang, X.-Y., et al. (2011). Bioaccumulation, maternal transfer and elimination of polybrominated diphenyl ethers in wild frogs. *Chemosphere*, 84(7), 972–8. doi:10.1016/j.chemosphere.2011.05.042
- 120.Liu, X., Ji, K., & Choi, K. (2012). Endocrine disruption potentials of organophosphate flame retardants and related mechanisms in H295R and MVLN cell lines and in zebrafish. *Aquatic toxicology (Amsterdam, Netherlands)*, 114-115, 173–81. doi:10.1016/j.aquatox.2012.02.019
- 121.Liu, X., Ji, K., Jo, A., Moon, H.-B., & Choi, K. (2013). Effects of TDCPP or TPP on gene transcriptions and hormones of HPG axis, and their consequences on reproduction in adult zebrafish (*Danio rerio*). *Aquatic toxicology (Amsterdam, Netherlands)*, 134-135, 104–11. doi:10.1016/j.aquatox.2013.03.013
- 122.Liu, Y., Li, J., Zhao, Y., Wen, S., Huang, F., & Wu, Y. (2011). Polybrominated diphenyl ethers (PBDEs) and indicator polychlorinated biphenyls (PCBs) in marine fish from four areas of China. *Chemosphere*, 83(2), 168–74. doi:10.1016/j.chemosphere.2010.12.045
- 123.Luo, X.-J., Zhang, X.-L., Chen, S.-J., & Mai, B.-X. (2010). Free and bound polybrominated diphenyl ethers and tetrabromobisphenol A in freshwater sediments. *Marine pollution bulletin*, 60(5), 718–24. doi:10.1016/j.marpolbul.2009.11.023
- 124.Ma, J., Qiu, X., Zhang, J., Duan, X., & Zhu, T. (2012). State of polybrominated diphenyl ethers in China: an overview. *Chemosphere*, 88(7), 769–78. doi:10.1016/j.chemosphere.2012.03.093
- 125.Ma, Y., Venier, M., & Hites, R. A. (2012). 2-Ethylhexyl tetrabromobenzoate and bis(2-ethylhexyl) tetrabromophthalate flame retardants in the Great Lakes atmosphere. *Environmental science & technology*, 46(1), 204–8. doi:10.1021/es203251f
- 126.Makris, K. C., & Snyder, S. A. (2010). Screening of pharmaceuticals and endocrine disrupting compounds in water supplies of Cyprus. *Water science and technology □: a journal of the International Association on Water Pollution Research*, 62(11), 2720–8. doi:10.2166/wst.2010.549
- 127.Marteinson, S. C., Bird, D. M., Shutt, J. L., Letcher, R. J., Ritchie, I. J., & Fernie, K. J. (2010). Multi-generational effects of polybrominated diphenylethers exposure: embryonic exposure of male American kestrels (*Falco sparverius*) to DE-71 alters reproductive success and behaviors. *Environmental toxicology and chemistry / SETAC*, 29(8), 1740–7. doi:10.1002/etc.200
- 128.Marteinson, S. C., Kimmins, S., Bird, D. M., Shutt, J. L., Letcher, R. J., Ritchie, I. J., & Fernie, K. J. (2011). Embryonic exposure to the polybrominated diphenyl ether mixture, DE-71, affects testes and circulating testosterone concentrations in adult American kestrels (*Falco sparverius*). *Toxicological sciences □: an official journal of the Society of Toxicology*, 121(1), 168–76. doi:10.1093/toxsci/kfr027
- 129.Martin, P. A., Mayne, G. J., Bursian, F. S. J., Tomy, G., Palace, V., Pekarik, C., & Smits, J. (2007). Immunotoxicity of the commercial polybrominated diphenyl ether mixture DE-71 in ranch mink (*Mustela vison*). *Environmental toxicology and chemistry / SETAC*, 26(5), 988–97. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17521147>
- 130.McClain, V., Stapleton, H. M., Tilton, F., & Gallagher, E. P. (2012). BDE 49 and developmental toxicity in zebrafish. *Comparative biochemistry and physiology. Toxicology & pharmacology □: CBP*, 155(2), 253–8. doi:10.1016/j.cbpc.2011.09.004
- 131.McGee, S. P., Cooper, E. M., Stapleton, H. M., & Volz, D. C. (2012). Early zebrafish embryogenesis is susceptible to developmental TDCPP exposure. *Environmental health perspectives*, 120(11), 1585–91. doi:10.1289/ehp.1205316
- 132.McGee, S. P., Konstantinov, A., Stapleton, H. M., & Volz, D. C. (2013a). Aryl phosphate esters within a major PentaBDE replacement product induce cardiotoxicity in developing zebrafish embryos: potential role of the aryl hydrocarbon receptor. *Toxicological sciences □: an official journal of the Society of Toxicology*, 133(1), 144–56. doi:10.1093/toxsci/kft020
- 133.McGee, S. P., Konstantinov, A., Stapleton, H. M., & Volz, D. C. (2013b). Aryl phosphate esters within a major PentaBDE replacement product induce cardiotoxicity in developing zebrafish embryos: potential role of the aryl

- hydrocarbon receptor. *Toxicological sciences* □: an official journal of the Society of Toxicology, 133(1), 144–56. doi:10.1093/toxsci/kft020
134. McKernan, M. A., Rattner, B. A., Hale, R. C., & Ottinger, M. A. (2009). Toxicity of polybrominated diphenyl ethers (DE-71) in chicken (*Gallus gallus*), mallard (*Anas platyrhynchos*), and American kestrel (*Falco sparverius*) embryos and hatchlings. *Environmental toxicology and chemistry / SETAC*, 28(5), 1007–17. doi:10.1897/08-318.1
135. McKinney, M. A., Dietz, R., Sonne, C., De Guise, S., Skirnisson, K., Karlsson, K., Steingrímsson, E., et al. (2011). Comparative hepatic microsomal biotransformation of selected PBDEs, including decabromodiphenyl ether, and decabromodiphenyl ethane flame retardants in Arctic marine-feeding mammals. *Environmental toxicology and chemistry / SETAC*, 30(7), 1506–14. doi:10.1002/etc.535
136. McKinney, M. A., Letcher, R. J., Aars, J., Born, E. W., Branigan, M., Dietz, R., Evans, T. J., Gabrielsen, G. W., Muir, D. C. G., et al. (2011). Regional contamination versus regional dietary differences: understanding geographic variation in brominated and chlorinated contaminant levels in polar bears. *Environmental science & technology*, 45(3), 896–902. doi:10.1021/es102781b
137. McKinney, M. A., Letcher, R. J., Aars, J., Born, E. W., Branigan, M., Dietz, R., Evans, T. J., Gabrielsen, G. W., Peacock, E., et al. (2011). Flame retardants and legacy contaminants in polar bears from Alaska, Canada, East Greenland and Svalbard, 2005–2008. *Environment international*, 37(2), 365–74. doi:10.1016/j.envint.2010.10.008
138. McKinney, M. A., Stirling, I., Lunn, N. J., Peacock, E., & Letcher, R. J. (2010). The role of diet on long-term concentration and pattern trends of brominated and chlorinated contaminants in western Hudson Bay polar bears, 1991–2007. *The Science of the total environment*, 408(24), 6210–22. doi:10.1016/j.scitotenv.2010.08.033
139. Meng, X.-Z., Blasius, M. E., Gossett, R. W., & Maruya, K. A. (2009). Polybrominated diphenyl ethers in pinnipeds stranded along the southern California coast. *Environmental pollution*, 157(10), 2731–6. doi:10.1016/j.envpol.2009.04.029
140. Meyer, T., Muir, D. C. G., Teixeira, C., Wang, X., Young, T., & Wania, F. (2012). Deposition of brominated flame retardants to the Devon Ice Cap, Nunavut, Canada. *Environmental science & technology*, 46(2), 826–33. doi:10.1021/es202900u
141. Mhadhbi, L., Fumega, J., Boumaiza, M., & Beiras, R. (2012). Acute toxicity of polybrominated diphenyl ethers (PBDEs) for turbot (*Psetta maxima*) early life stages (ELS). *Environmental science and pollution research international*, 19(3), 708–17. doi:10.1007/s11356-011-0602-5
142. Miljeteig, C., Gabrielsen, G. W., Strøm, H., Gavrilo, M. V., Lie, E., & Jenssen, B. M. (2012). Eggshell thinning and decreased concentrations of vitamin E are associated with contaminants in eggs of ivory gulls. *The Science of the total environment*, 431, 92–9. doi:10.1016/j.scitotenv.2012.05.018
143. Mo, L., Wu, J.-P., Luo, X.-J., Zou, F.-S., & Mai, B.-X. (2012). Bioaccumulation of polybrominated diphenyl ethers, decabromodiphenyl ethane, and 1,2-bis(2,4,6-tribromophenoxy) ethane flame retardants in kingfishers (*Alcedo atthis*) from an electronic waste-recycling site in South China. *Environmental toxicology and chemistry / SETAC*, 31(9), 2153–8. doi:10.1002/etc.1929
144. Möller, A., Sturm, R., Xie, Z., Cai, M., He, J., & Ebinghaus, R. (2012). Organophosphorus flame retardants and plasticizers in airborne particles over the Northern Pacific and Indian Ocean toward the Polar Regions: evidence for global occurrence. *Environmental science & technology*, 46(6), 3127–34. doi:10.1021/es204272v
145. Möller, A., Xie, Z., Caba, A., Sturm, R., & Ebinghaus, R. (2011). Organophosphorus flame retardants and plasticizers in the atmosphere of the North Sea. *Environmental pollution (Barking, Essex □: 1987)*, 159(12), 3660–5. doi:10.1016/j.envpol.2011.07.022
146. Möller, A., Xie, Z., Cai, M., Sturm, R., & Ebinghaus, R. (2012). Brominated flame retardants and dechlorane plus in the marine atmosphere from Southeast Asia toward Antarctica. *Environmental science & technology*, 46(6), 3141–8. doi:10.1021/es300138q

147. Möller, A., Xie, Z., Cai, M., Zhong, G., Huang, P., Cai, M., Sturm, R., et al. (2011). Polybrominated diphenyl ethers vs alternate brominated flame retardants and Dechloranes from East Asia to the Arctic. *Environmental science & technology*, 45(16), 6793–9. doi:10.1021/es201850n
148. Möller, A., Xie, Z., Sturm, R., & Ebinghaus, R. (2011). Polybrominated diphenyl ethers (PBDEs) and alternative brominated flame retardants in air and seawater of the European Arctic. *Environmental pollution (Barking, Essex□: 1987)*, 159(6), 1577–83. doi:10.1016/j.envpol.2011.02.054
149. Montie, E. W., Letcher, R. J., Reddy, C. M., Moore, M. J., Rubinstein, B., & Hahn, M. E. (2010). Brominated flame retardants and organochlorine contaminants in winter flounder, harp and hooded seals, and North Atlantic right whales from the Northwest Atlantic Ocean. *Marine pollution bulletin*, 60(8), 1160–9. doi:10.1016/j.marpolbul.2010.04.002
150. Montory, M., Habit, E., Fernandez, P., Grimalt, J. O., & Barra, R. (2012). Polybrominated diphenyl ether levels in wild and farmed Chilean salmon and preliminary flow data for commercial transport. *Journal of environmental sciences (China)*, 24(2), 221–7. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22655380>
151. Morrissey, C. A., Pollet, I. L., Ormerod, S. J., & Elliott, J. E. (2012). American dippers indicate contaminant biotransport by Pacific salmon. *Environmental science & technology*, 46(2), 1153–62. doi:10.1021/es2028058
152. Muirhead, E. K., Skillman, A. D., Hook, S. E., & Schultz, I. R. (2006). Oral exposure of PBDE-47 in fish: toxicokinetics and reproductive effects in Japanese Medaka (*Oryzias latipes*) and fathead minnows (*Pimephales promelas*). *Environmental science & technology*, 40(2), 523–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16468398>
153. Mukherjee, J., Po, B. H. K., Chiu, J. M. Y., Wu, R. S. S., Qian, P.-Y., & Thiagarajan, V. (2013). Polybrominated diphenyl ethers do not affect metamorphosis but alter the proteome of the invasive slipper limpet *Crepidula onyx*. *Marine pollution bulletin*. doi:10.1016/j.marpolbul.2013.05.009
154. Muñoz-Arnanz, J., Sáez, M., Aguirre, J. I., Hidalgo, F., Baos, R., Pacepavicius, G., Alaee, M., et al. (2011). Predominance of BDE-209 and other higher brominated diphenyl ethers in eggs of white stork (*Ciconia ciconia*) colonies from Spain. *Environment international*, 37(3), 572–6. doi:10.1016/j.envint.2010.11.013
155. Munsch, C., Héas-Moisan, K., Tixier, C., Boulesteix, L., & Morin, J. (2011). Classic and novel brominated flame retardants (BFRs) in common sole (*Solea solea* L.) from main nursery zones along the French coasts. *The Science of the total environment*, 409(21), 4618–27. doi:10.1016/j.scitotenv.2011.07.021
156. Newsome, S. D., Park, J.-S., Henry, B. W., Holden, A., Fogel, M. L., Linthicum, J., Chu, V., et al. (2010). Polybrominated diphenyl ether (PBDE) levels in peregrine falcon (*Falco peregrinus*) eggs from California correlate with diet and human population density. *Environmental science & technology*, 44(13), 5248–55. doi:10.1021/es100658e
157. Nordlöf, U., Helander, B., Bignert, A., & Asplund, L. (2010). Levels of brominated flame retardants and methoxylated polybrominated diphenyl ethers in eggs of white-tailed sea eagles breeding in different regions of Sweden. *The Science of the total environment*, 409(1), 238–46. doi:10.1016/j.scitotenv.2010.09.042
158. Nøst, T. H., Helgason, L. B., Harju, M., Heimstad, E. S., Gabrielsen, G. W., & Jenssen, B. M. (2012). Halogenated organic contaminants and their correlations with circulating thyroid hormones in developing Arctic seabirds. *The Science of the total environment*, 414, 248–56. doi:10.1016/j.scitotenv.2011.11.051
159. Noyes, P. D., Hinton, D. E., & Stapleton, H. M. (2011). Accumulation and debromination of decabromodiphenyl ether (BDE-209) in juvenile fathead minnows (*Pimephales promelas*) induces thyroid disruption and liver alterations. *Toxicological sciences□: an official journal of the Society of Toxicology*, 122(2), 265–74. doi:10.1093/toxsci/kfr105
160. Nyholm, J. R., Asamoah, R. K., Van der Wal, L., Danielsson, C., & Andersson, P. L. (2010). Accumulation of polybrominated diphenyl ethers, hexabromobenzene, and 1,2-dibromo-4-(1,2-dibromoethyl)cyclohexane in earthworm (*Eisenia fetida*). Effects of soil type and aging. *Environmental science & technology*, 44(23), 9189–94. doi:10.1021/es1023288

- 161.Nyholm, J. R., Grbic, R., Arp, H. P. H., Moskeland, T., & Andersson, P. L. (2013). Environmental occurrence of emerging and legacy brominated flame retardants near suspected sources in Norway. *The Science of the total environment*, 443, 307–14. doi:10.1016/j.scitotenv.2012.10.081
- 162.Odusanya, D. O., Okonkwo, J. O., & Botha, B. (2009). Polybrominated diphenyl ethers (PBDEs) in leachates from selected landfill sites in South Africa. *Waste management*, 29(1), 96–102. doi:10.1016/j.wasman.2008.02.011
- 163.Oliaei, F., King, P., & Phillips, L. (2002). Occurrence and concentrations of polybrominated diphenyl ethers (PBDEs) in Minnesota environment. *Organohalogen Compounds*, 58, 185–188. Retrieved from <http://www.dioxin20xx.org/pdfs/2002/02-419.pdf>
- 164.Olshansky, Y., Polubesova, T., Vetter, W., & Chefetz, B. (2011). Sorption-desorption behavior of polybrominated diphenyl ethers in soils. *Environmental pollution (Barking, Essex, England)*, 159(10), 2375–9. doi:10.1016/j.envpol.2011.07.003
- 165.Osako, M., Kim, Y.-J., & Sakai, S. (2004). Leaching of brominated flame retardants in leachate from landfills in Japan. *Chemosphere*, 57(10), 1571–9. doi:10.1016/j.chemosphere.2004.08.076
- 166.Pan, X., Tang, J., Li, J., Zhong, G., Chen, Y., & Zhang, G. (2011). Polybrominated diphenyl ethers (PBDEs) in the riverine and marine sediments of the Laizhou Bay area, North China. *Journal of environmental monitoring* □: JEM, 13(4), 886–93. doi:10.1039/c1em10169b
- 167.Parolini, M., & Binelli, A. (2012). Cyto-genotoxic effects induced by three brominated diphenyl ether congeners on the freshwater mussel *Dreissena polymorpha*. *Ecotoxicology and environmental safety*, 79, 247–55. doi:10.1016/j.ecoenv.2012.01.008
- 168.Petreas, M., & Oros, D. (2009). Polybrominated diphenyl ethers in California wastestreams. *Chemosphere*, 74(7), 996–1001. doi:10.1016/j.chemosphere.2008.09.093
- 169.Pujolar, J. M., Milan, M., Marino, I. A. M., Capoccioni, F., Ciccotti, E., Belpaire, C., Covaci, A., et al. (2013). Detecting genome-wide gene transcription profiles associated with high pollution burden in the critically endangered European eel. *Aquatic toxicology (Amsterdam, Netherlands)*, 132-133, 157–64. doi:10.1016/j.aquatox.2013.02.012
- 170.Qiu, X., Zhu, T., & Hu, J. (2010). Polybrominated diphenyl ethers (PBDEs) and other flame retardants in the atmosphere and water from Taihu Lake, East China. *Chemosphere*, 80(10), 1207–12. doi:10.1016/j.chemosphere.2010.06.013
- 171.Raach, M., Lebeuf, M., & Pelletier, E. (2011). PBDEs and PCBs in the liver of the St Lawrence Estuary beluga (*Delphinapterus leucas*): a comparison of levels and temporal trends with the blubber. *Journal of environmental monitoring* □: JEM. doi:10.1039/c0em00310g
- 172.Rahman, F., Langford, K. H., Scrimshaw, M. D., & Lester, J. N. (2001). Polybrominated diphenyl ether (PBDE) flame retardants. *The Science of the total environment*, 275(1-3), 1–17. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11482396>
- 173.Rawn, D. F. K., Sadler, A., Quade, S. C., Sun, W.-F., Lau, B. P.-Y., Kosarac, I., Hayward, S., et al. (2011). Brominated flame retardants in Canadian chicken egg yolks. *Food additives & contaminants. Part A, Chemistry, analysis, control, exposure & risk assessment*, 28(6), 807–15. doi:10.1080/19440049.2010.545443
- 174.Rayne, S., Ikonomou, M. G., Ross, P. S., Ellis, G. M., & Barrett-Lennard, L. G. (2004). PBDEs, PBBs, and PCNs in three communities of free-ranging killer whales (*Orcinus orca*) from the northeastern Pacific Ocean. *Environmental science & technology*, 38(16), 4293–9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15382855>
- 175.REEMTSMA, T., QUINTANA, J., RODIL, R., GARCIA LOPEZ, M., & RODRIGUEZ, I. (2008). Organophosphorus flame retardants and plasticizers in water and air I. Occurrence and fate. *Trends in Analytical Chemistry*, 27(9), 727–737. doi:10.1016/j.trac.2008.07.002
- 176.Regnery, J., Püttmann, W., Merz, C., & Berthold, G. (2011). Occurrence and distribution of organophosphorus flame retardants and plasticizers in anthropogenically affected groundwater. *Journal of environmental monitoring* □: JEM, 13(2), 347–54. doi:10.1039/c0em00419g

177. Regnery, Julia, & Püttmann, W. (2010a). Seasonal fluctuations of organophosphate concentrations in precipitation and storm water runoff. *Chemosphere*, 78(8), 958–64. doi:10.1016/j.chemosphere.2009.12.027
178. Regnery, Julia, & Püttmann, W. (2010b). Occurrence and fate of organophosphorus flame retardants and plasticizers in urban and remote surface waters in Germany. *Water research*, 44(14), 4097–104. doi:10.1016/j.watres.2010.05.024
179. Ricklund, N., Kierkegaard, A., & McLachlan, M. S. (2008). An international survey of decabromodiphenyl ethane (deBDethane) and decabromodiphenyl ether (decaBDE) in sewage sludge samples. *Chemosphere*, 73(11), 1799–804. doi:10.1016/j.chemosphere.2008.08.047
180. Ricklund, N., Kierkegaard, A., & McLachlan, M. S. (2010). Levels and potential sources of decabromodiphenyl ethane (DBDPE) and decabromodiphenyl ether (DecaBDE) in lake and marine sediments in Sweden. *Environmental science & technology*, 44(6), 1987–91. doi:10.1021/es903701q
181. Rodil, R., Quintana, J. B., Concha-Graña, E., López-Mahía, P., Muniategui-Lorenzo, S., & Prada-Rodríguez, D. (2012). Emerging pollutants in sewage, surface and drinking water in Galicia (NW Spain). *Chemosphere*, 86(10), 1040–9. doi:10.1016/j.chemosphere.2011.11.053
182. Ross, P. S. (2006). Fireproof killer whales (Orcinus orca): flame-retardant chemicals and the conservation imperative in the charismatic icon of British Columbia, Canada. *Canadian Journal of Fisheries and Aquatic Sciences*, 63(1), 224–234. doi:10.1139/f05-244
183. Ross, P. S., Couillard, C. M., Ikonomou, M. G., Johannessen, S. C., Lebeuf, M., Macdonald, R. W., & Tomy, G. T. (2009). Large and growing environmental reservoirs of Deca-BDE present an emerging health risk for fish and marine mammals. *Marine pollution bulletin*, 58(1), 7–10. doi:10.1016/j.marpolbul.2008.09.002
184. Routti, H., Helgason, L. B., Arukwe, A., Wolkers, H., Heimstad, E. S., Harju, M., Berg, V., et al. (2013). Effect of reduced food intake on toxicokinetics of halogenated organic contaminants in herring gull (*Larus argentatus*) chicks. *Environmental toxicology and chemistry / SETAC*, 32(1), 156–64. doi:10.1002/etc.2026
185. Sagerup, K., Herzke, D., Harju, M., Evensen, A., Christensen, G., Routti, H., Fuglei, E., et al. (2010). New brominated flame retardants in Arctic biota (TA-2630-2010). *Norwegian Climate and Pollution Agency (Klif)*. Retrieved from <http://www.klif.no/publikasjoner/2630/ta2630.pdf>
186. Salamova, A., & Hites, R. A. (2011). Discontinued and alternative brominated flame retardants in the atmosphere and precipitation from the great lakes basin. *Environmental science & technology*, 45(20), 8698–706. doi:10.1021/es2020378
187. Salamova, A., & Hites, R. A. (2013). Brominated and chlorinated flame retardants in tree bark from around the globe. *Environmental science & technology*, 47(1), 349–54. doi:10.1021/es303393z
188. Shaw, S. D., Berger, M. L., Brenner, D., Kannan, K., Lohmann, N., & Päpke, O. (2009). Bioaccumulation of polybrominated diphenyl ethers and hexabromocyclododecane in the northwest Atlantic marine food web. *The Science of the total environment*, 407(10), 3323–9. doi:10.1016/j.scitotenv.2009.02.018
189. Shaw, S. D., Berger, M. L., Weijs, L., & Covaci, A. (2012). Tissue-specific accumulation of polybrominated diphenyl ethers (PBDEs) including Deca-BDE and hexabromocyclododecanes (HBCDs) in harbor seals from the northwest Atlantic. *Environment international*, 44, 1–6. doi:10.1016/j.envint.2012.01.001
190. Shaw, S. D., Brenner, D., Berger, M. L., Fang, F., Hong, C.-S., Addink, R., & Hilker, D. (2008). Bioaccumulation of polybrominated diphenyl ethers in harbor seals from the northwest Atlantic. *Chemosphere*, 73(11), 1773–80. doi:10.1016/j.chemosphere.2008.09.016
191. Shaw, S. D., & Kannan, K. (n.d.). Polybrominated diphenyl ethers in marine ecosystems of the American continents: foresight from current knowledge. *Reviews on environmental health*, 24(3), 157–229. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19891120>
192. She, J., Petreas, M., Winkler, J., Visita, P., McKinney, M., & Kopec, D. (2002). PBDEs in the San Francisco Bay Area: measurements in harbor seal blubber and human breast adipose tissue. *Chemosphere*, 46(5), 697–707. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11999793>

193. Søfteland, L., Petersen, K., Stavrum, A.-K., Wu, T., & Olsvik, P. A. (2011). Hepatic in vitro toxicity assessment of PBDE congeners BDE47, BDE153 and BDE154 in Atlantic salmon (*Salmo salar* L.). *Aquatic toxicology (Amsterdam, Netherlands)*, *105*(3-4), 246–63. doi:10.1016/j.aquatox.2011.03.012
194. Sørmo, E. G., Lie, E., Ruus, A., Gaustad, H., Skaare, J. U., & Jenssen, B. M. (2011). Trophic level determines levels of brominated flame-retardants in coastal herring gulls. *Ecotoxicology and environmental safety*, *74*(7), 2091–8. doi:10.1016/j.ecoenv.2011.06.012
195. Stapleton, H M, Dodder, N. G., Kucklick, J. R., Reddy, C. M., Schantz, M. M., Becker, P. R., Gulland, F., et al. (2006). Determination of HBCD, PBDEs and MeO-BDEs in California sea lions (*Zalophus californianus*) stranded between 1993 and 2003. *Marine pollution bulletin*, *52*(5), 522–31. doi:10.1016/j.marpolbul.2005.09.045
196. Stapleton, Heather M, Alaee, M., Letcher, R. J., & Baker, J. E. (2004). Debromination of the flame retardant decabromodiphenyl ether by juvenile carp (*Cyprinus carpio*) following dietary exposure. *Environmental science & technology*, *38*(1), 112–9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14740725>
197. Su, G., Zhang, X., Liu, H., Giesy, J. P., Lam, M. H. W., Lam, P. K. S., Siddiqui, M. A., et al. (2012). Toxicogenomic mechanisms of 6-HO-BDE-47, 6-MeO-BDE-47, and BDE-47 in *E. coli*. *Environmental science & technology*, *46*(2), 1185–91. doi:10.1021/es203212w
198. Sühring, R., Möller, A., Freese, M., Pohlmann, J.-D., Wolschke, H., Sturm, R., Xie, Z., et al. (2013). Brominated flame retardants and dechlorananes in eels from German Rivers. *Chemosphere*, *90*(1), 118–24. doi:10.1016/j.chemosphere.2012.08.016
199. Sullivan, K. M., Bird, D. M., Ritchie, J. I., Shutt, J. L., Letcher, R. J., & Fernie, K. J. (2010). Changes in plasma retinol of American kestrels (*Falco sparverius*) in response to dietary or in ovo exposure to environmentally relevant concentrations of a penta-brominated diphenyl ether mixture, DE-71. *Journal of toxicology and environmental health. Part A*, *73*(23), 1645–54. doi:10.1080/15287394.2010.501720
200. Sun, Y., Luo, X., Mo, L., Zhang, Q., Wu, J., Chen, S., Zou, F., et al. (2012). Brominated flame retardants in three terrestrial passerine birds from South China: geographical pattern and implication for potential sources. *Environmental pollution (Barking, Essex, England : 1987)*, *162*, 381–8. doi:10.1016/j.envpol.2011.12.013
201. Sundkvist, A. M., Olofsson, U., & Haglund, P. (2010). Organophosphorus flame retardants and plasticizers in marine and fresh water biota and in human milk. *Journal of environmental monitoring (London : 1997) : JEM*, *12*(4), 943–51. doi:10.1039/b921910b
202. Suutari, A., Ruokojärvi, P., Kiviranta, H., Verta, M., Korhonen, M., Nieminen, M., & Laaksonen, S. (2011). Polychlorinated dibenzo-p-dioxins (PCDDs), dibenzofurans (PCDFs), polychlorinated biphenyls (PCBs), and polybrominated diphenyl ethers (PBDEs) in Finnish semi-domesticated reindeer (*Rangifer tarandus tarandus* L.). *Environment international*, *37*(2), 335–41. doi:10.1016/j.envint.2010.10.003
203. Takahashi, S., Oshihoi, T., Ramu, K., Isobe, T., Ohmori, K., Kubodera, T., & Tanabe, S. (2010). Organohalogen compounds in deep-sea fishes from the western North Pacific, off-Tohoku, Japan: Contamination status and bioaccumulation profiles. *Marine pollution bulletin*, *60*(2), 187–96. doi:10.1016/j.marpolbul.2009.09.027
204. Tanabe, S., Ramu, K., Isobe, T., & Takahashi, S. (2008). Brominated flame retardants in the environment of Asia-Pacific: an overview of spatial and temporal trends. *Journal of environmental monitoring (London : 1997) : JEM*, *10*(2), 188–97. doi:10.1039/b709928b
205. Tapie, N., Le Menach, K., Pasquaud, S., Elie, P., Devier, M. H., & Budzinski, H. (2011). PBDE and PCB contamination of eels from the Gironde estuary: from glass eels to silver eels. *Chemosphere*, *83*(2), 175–85. doi:10.1016/j.chemosphere.2010.12.044
206. Tian, M., Chen, S.-J., Luo, Y., Wang, J., Zhu, Z.-C., Luo, X.-J., & Mai, B.-X. (2013). Air-plant exchange of brominated flame retardants at a rural site: Influencing factor, interspecies difference, and forest scavenging. *Environmental toxicology and chemistry / SETAC*, *32*(6), 1248–53. doi:10.1002/etc.2195

- 207.Timme-Laragy, A. R., Levin, E. D., & Di Giulio, R. T. (2006). Developmental and behavioral effects of embryonic exposure to the polybrominated diphenylether mixture DE-71 in the killifish (*Fundulus heteroclitus*). *Chemosphere*, 62(7), 1097–104. doi:10.1016/j.chemosphere.2005.05.037
- 208.Tomy, G. T., Palace, V. P., Halldorson, T., Braekevelt, E., Danell, R., Wautier, K., Evans, B., et al. (2004). Bioaccumulation, biotransformation, and biochemical effects of brominated diphenyl ethers in juvenile lake trout (*Salvelinus namaycush*). *Environmental science & technology*, 38(5), 1496–504. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15046352>
- 209.Torres, L., Orazio, C. E., Peterman, P. H., & Patiño, R. (2013). Effects of dietary exposure to brominated flame retardant BDE-47 on thyroid condition, gonadal development and growth of zebrafish. *Fish physiology and biochemistry*. doi:10.1007/s10695-012-9768-0
- 210.Ueno, D., Isobe, T., Ramu, K., Tanabe, S., Alaee, M., Marvin, C., Inoue, K., et al. (2010). Spatial distribution of hexabromocyclododecanes (HBCDs), polybrominated diphenyl ethers (PBDEs) and organochlorines in bivalves from Japanese coastal waters. *Chemosphere*, 78(10), 1213–9. doi:10.1016/j.chemosphere.2009.12.058
- 211.Van den Steen, E., Eens, M., Covaci, A., Durtu, A. C., Jaspers, V. L. B., Neels, H., & Pinxten, R. (2009). An exposure study with polybrominated diphenyl ethers (PBDEs) in female European starlings (*Sturnus vulgaris*): toxicokinetics and reproductive effects. *Environmental pollution*, 157(2), 430–6. doi:10.1016/j.envpol.2008.09.031
- 212.Van der Veen, I., & De Boer, J. (2012). Phosphorus flame retardants: properties, production, environmental occurrence, toxicity and analysis. *Chemosphere*, 88(10), 1119–53. doi:10.1016/j.chemosphere.2012.03.067
- 213.Van Schmidt, N. D., Cary, T. L., Ortiz-Santaliestra, M. E., & Karasov, W. H. (2012). Effects of chronic polybrominated diphenyl ether exposure on gonadal development in the northern leopard frog, *Rana pipiens*. *Environmental toxicology and chemistry / SETAC*, 31(2), 347–54. doi:10.1002/etc.1702
- 214.Vane, C. H., Ma, Y.-J., Chen, S.-J., & Mai, B.-X. (2010). Increasing polybrominated diphenyl ether (PBDE) contamination in sediment cores from the inner Clyde Estuary, UK. *Environmental geochemistry and health*, 32(1), 13–21. doi:10.1007/s10653-009-9261-6
- 215.Venier, M., & Hites, R. A. (2011). Flame retardants in the serum of pet dogs and in their food. *Environmental science & technology*, 45(10), 4602–8. doi:10.1021/es1043529
- 216.Venier, M., Wierda, M., Bowerman, W. W., & Hites, R. A. (2010). Flame retardants and organochlorine pollutants in bald eagle plasma from the Great Lakes region. *Chemosphere*, 80(10), 1234–40. doi:10.1016/j.chemosphere.2010.05.043
- 217.Verboven, N., Verreault, J., Letcher, R. J., Gabrielsen, G. W., & Evans, N. P. (2010). Adrenocortical function of Arctic-breeding glaucous gulls in relation to persistent organic pollutants. *General and comparative endocrinology*, 166(1), 25–32. doi:10.1016/j.ygcen.2009.11.013
- 218.Verreault, J., Gebbink, W. A., Gauthier, L. T., Gabrielsen, G. W., & Letcher, R. J. (2007). Brominated flame retardants in glaucous gulls from the Norwegian Arctic: more than just an issue of polybrominated diphenyl ethers. *Environmental science & technology*, 41(14), 4925–31. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17711204>
- 219.Viganò, L., Roscioli, C., & Guzzella, L. (2011). Decabromodiphenyl ether (BDE-209) enters the food web of the River Po and is metabolically debrominated in resident cyprinid fishes. *The Science of the total environment*, 409(23), 4966–72. doi:10.1016/j.scitotenv.2011.07.062
- 220.Villanger, G. D., Lydersen, C., Kovacs, K. M., Lie, E., Skaare, J. U., & Jenssen, B. M. (2011). Disruptive effects of persistent organohalogen contaminants on thyroid function in white whales (*Delphinapterus leucas*) from Svalbard. *The Science of the total environment*, 409(13), 2511–24. doi:10.1016/j.scitotenv.2011.03.014
- 221.Vrkoslavová, J., Demnerová, K., Macková, M., Zemanová, T., Macek, T., Hajslová, J., Pulkrabová, J., et al. (2010). Absorption and translocation of polybrominated diphenyl ethers (PBDEs) by plants from contaminated sewage sludge. *Chemosphere*, 81(3), 381–6. doi:10.1016/j.chemosphere.2010.07.010

222. Waaijers, S. L., Kong, D., Hendriks, H. S., De Wit, C. A., Cousins, I. T., Westerink, R. H. S., Leonards, P. E. G., et al. (2013). Persistence, bioaccumulation, and toxicity of halogen-free flame retardants. *Reviews of environmental contamination and toxicology*, 222, 1–71. doi:10.1007/978-1-4614-4717-7_1
223. Wang, C., Li, W., Chen, J., Wang, H., Li, T., Shen, G., Shen, H., et al. (2012). Summer atmospheric polybrominated diphenyl ethers in urban and rural areas of northern China. *Environmental pollution (Barking, Essex)*, 171, 234–40. doi:10.1016/j.envpol.2012.07.041
224. Wang, D.-G., Alaee, M., Sverko, E., Li, Y.-F., Reiner, E. J., & Shen, L. (2011). Analysis and occurrence of emerging chlorinated and brominated flame retardants in surficial sediment of the Dalian coastal area in China. *Journal of environmental monitoring*, 13(11), 3104–10. doi:10.1039/c1em10241a
225. Wang, J., Kliks, M. M., Jun, S., & Li, Q. X. (2010). Residues of polybrominated diphenyl ethers in honeys from different geographic regions. *Journal of agricultural and food chemistry*, 58(6), 3495–501. doi:10.1021/jf904490g
226. Wang, L.-C., Hsi, H.-C., Wang, Y.-F., Lin, S.-L., & Chang-Chien, G.-P. (2010). Distribution of polybrominated diphenyl ethers (PBDEs) and polybrominated dibenzo-p-dioxins and dibenzofurans (PBDD/Fs) in municipal solid waste incinerators. *Environmental pollution*, 158(5), 1595–602. doi:10.1016/j.envpol.2009.12.016
227. Wang, S., Zhang, S., Huang, H., & Christie, P. (2011). Behavior of decabromodiphenyl ether (BDE-209) in soil: effects of rhizosphere and mycorrhizal colonization of ryegrass roots. *Environmental pollution (Barking, Essex)*, 159(3), 749–53. doi:10.1016/j.envpol.2010.11.035
228. Weijs, L., Das, K., Neels, H., Blust, R., & Covaci, A. (2010). Occurrence of anthropogenic and naturally-produced organohalogenated compounds in tissues of Black Sea harbour porpoises. *Marine pollution bulletin*, 60(5), 725–31. doi:10.1016/j.marpolbul.2009.11.022
229. Wiseman, S. B., Wan, Y., Chang, H., Zhang, X., Hecker, M., Jones, P. D., & Giesy, J. P. (2011). Polybrominated diphenyl ethers and their hydroxylated/methoxylated analogs: environmental sources, metabolic relationships, and relative toxicities. *Marine pollution bulletin*, 63(5–12), 179–88. doi:10.1016/j.marpolbul.2011.02.008
230. Wong, F., Kurt-Karakus, P., & Bidleman, T. F. (2012). Fate of brominated flame retardants and organochlorine pesticides in urban soil: volatility and degradation. *Environmental science & technology*, 46(5), 2668–74. doi:10.1021/es203287x
231. Wu, J., Zhang, Y., Luo, X., She, Y., Yu, L., Chen, S., & Mai, B. (2012). A review of polybrominated diphenyl ethers and alternative brominated flame retardants in wildlife from China: levels, trends, and bioaccumulation characteristics. *Journal of environmental sciences (China)*, 24(2), 183–94. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22655375>
232. Xiao, H., Shen, L., Su, Y., Barresi, E., Dejong, M., Hung, H., Lei, Y.-D., et al. (2012). Atmospheric concentrations of halogenated flame retardants at two remote locations: the Canadian High Arctic and the Tibetan Plateau. *Environmental pollution (Barking, Essex)*, 161, 154–61. doi:10.1016/j.envpol.2011.09.041
233. Xie, Z., Möller, A., Ahrens, L., Sturm, R., & Ebinghaus, R. (2011). Brominated flame retardants in seawater and atmosphere of the Atlantic and the Southern Ocean. *Environmental science & technology*, 45(5), 1820–6. doi:10.1021/es103803t
234. Xu, Y., Zhang, G., Li, J., Liu, X., & Li, X. (2011). Atmospheric polybrominated diphenyl ethers (PBDEs) and Pb isotopes at a remote site in Southwestern China: implications for monsoon-associated transport. *The Science of the total environment*, 409(21), 4564–71. doi:10.1016/j.scitotenv.2011.07.035
235. Yogui, G. T., Santos, M. C. O., Bertozzi, C. P., Sericano, J. L., & Montone, R. C. (2011). PBDEs in the blubber of marine mammals from coastal areas of São Paulo, Brazil, southwestern Atlantic. *Marine pollution bulletin*, 62(12), 2666–70. doi:10.1016/j.marpolbul.2011.09.024
236. Yu, L., Deng, J., Shi, X., Liu, C., Yu, K., & Zhou, B. (2010). Exposure to DE-71 alters thyroid hormone levels and gene transcription in the hypothalamic-pituitary-thyroid axis of zebrafish larvae. *Aquatic toxicology (Amsterdam, Netherlands)*, 97(3), 226–33. doi:10.1016/j.aquatox.2009.10.022

- 237.Yu, L., Lam, J. C. W., Guo, Y., Wu, R. S. S., Lam, P. K. S., & Zhou, B. (2011). Parental transfer of polybrominated diphenyl ethers (PBDEs) and thyroid endocrine disruption in zebrafish. *Environmental science & technology*, 45(24), 10652–9. doi:10.1021/es2026592
- 238.Yu, L.-H., Luo, X.-J., Wu, J.-P., Liu, L.-Y., Song, J., Sun, Q.-H., Zhang, X.-L., et al. (2011). Biomagnification of higher brominated PBDE congeners in an urban terrestrial food web in north China based on field observation of prey deliveries. *Environmental science & technology*, 45(12), 5125–31. doi:10.1021/es200030z
- 239.Yu, Z., Liao, R., Li, H., Mo, L., Zeng, X., Sheng, G., & Fu, J. (2011). Particle-bound Dechlorane Plus and polybrominated diphenyl ethers in ambient air around Shanghai, China. *Environmental pollution (Barking, Essex)*, 159(10), 2982–8. doi:10.1016/j.envpol.2011.04.024
- 240.Zhang, S., Bursian, S. J., Martin, P., Chan, H. M., & Martin, J. (2008). Dietary accumulation, disposition, and metabolism of technical pentabrominated diphenyl ether (DE-71) in pregnant mink (*mustela vision*) and their offspring. *Environmental toxicology and chemistry / SETAC*, 1. doi:10.1897/07-487
- 241.Zhao, X., Zhang, H., Ni, Y., Lu, X., Zhang, X., Su, F., Fan, J., et al. (2011). Polybrominated diphenyl ethers in sediments of the Daliao River Estuary, China: levels, distribution and their influencing factors. *Chemosphere*, 82(9), 1262–7. doi:10.1016/j.chemosphere.2010.12.032
- 242.Zhu, N., Li, A., Wang, T., Wang, P., Qu, G., Ruan, T., Fu, J., et al. (2012). Tris(2,3-dibromopropyl) isocyanurate, hexabromocyclododecanes, and polybrominated diphenyl ethers in mollusks from Chinese Bohai Sea. *Environmental science & technology*, 46(13), 7174–81. doi:10.1021/es300776f
- 243.Zimmer, K. E., Montaño, M., Olsaker, I., Dahl, E., Berg, V., Karlsson, C., Murk, A. J., et al. (2011). In vitro steroidogenic effects of mixtures of persistent organic pollutants (POPs) extracted from burbot (*Lota lota*) caught in two Norwegian lakes. *The Science of the total environment*, 409(11), 2040–8. doi:10.1016/j.scitotenv.2011.01.055